

HYPOGONADOTROPIC HYPOGONADISM (KALLMANN)

PANEL DG-4.1.0 (58 GENES)

Gene	Twist X2 covered 10x	Twist X2 covered 20x	srWGS covered 10x	srWGS covered 15x	srWGS covered 20x	Associated Phenotype description and OMIM disease ID
ADCY3	100%	100%	100%	99.9%	99.2%	{Obesity, susceptibility to, BMIQ19}, 617885
ANOS1	100%	99.7%	98.5%	88.6%	71.5%	Hypogonadotropic hypogonadism 1 with or without anosmia (Kallmann syndrome 1), 308700
ARHGAP35	100%	100%	100%	99.9%	99.1%	
AXL	100%	100%	100%	99.9%	98.7%	
CCDC141	100%	100%	100%	100%	99.8%	
CHD7	100%	100%	100%	100%	99.4%	Hypogonadotropic hypogonadism 5 with or without anosmia, 612370;CHARGE syndrome, 214800
CNGA2	99.9%	98.7%	97.5%	84.7%	62.9%	

CUL4B	96.7%	96.5%	99.2%	91.3%	72.6%	Intellectual developmental disorder, X-linked syndromic, Cabezas type, 300354
DCAF17	100%	100%	100%	100%	99.4%	Woodhouse-Sakati syndrome, 241080
DCC	100%	100%	100%	99.9%	99.2%	Mirror movements 1 and/or agenesis of the corpus callosum, 157600;Esophageal carcinoma, somatic, 133239;Colorectal cancer, somatic, 114500;Gaze palsy, familial horizontal, with progressive scoliosis, 2, 617542
DUSP6	100%	100%	100%	100%	99.5%	Hypogonadotropic hypogonadism 19 with or without anosmia, 615269
FEZF1	100%	100%	100%	99.9%	99.2%	Hypogonadotropic hypogonadism 22, with or without anosmia, 616030
FGF17	100%	100%	100%	100%	99.2%	Hypogonadotropic hypogonadism 20 with or without anosmia, 615270

FGF8	100%	100%	100%	99.5%	97.3%	Hypogonadotropic hypogonadism 6 with or without anosmia, 612702
FGFR1	99.8%	98.9%	100%	100%	99.1%	Pfeiffer syndrome, 101600;Hypogonadotropic hypogonadism 2 with or without anosmia, 147950;Jackson-Weiss syndrome, 123150;Hartsfield syndrome, 615465;Trigonocephaly 1, 190440;Osteoglophonic dysplasia, 166250;Encephalocraniocutaneous lipomatosis, somatic mosaic, 613001
FLRT3	100%	100%	100%	100%	99.8%	Hypogonadotropic hypogonadism 21 with anosmia, 615271
FSHB	100%	100%	100%	100%	99.9%	Hypogonadotropic hypogonadism 24 without anosmia, 229070
GLI2	100%	100%	100%	99.9%	98.7%	Culler-Jones syndrome, 615849;Holoprosencephaly 9, 610829

GNRH1	100%	100%	100%	100%	100%	?Hypogonadotropic hypogonadism 12 with or without anosmia, 614841
GNRHR	100%	100%	100%	100%	99.7%	Hypogonadotropic hypogonadism 7 without anosmia, 146110
HESX1	100%	100%	100%	100%	99.5%	Pituitary hormone deficiency, combined, 5, 182230;Septo-optic dysplasia, 182230;Growth hormone deficiency with pituitary anomalies, 182230
HS6ST1	100%	100%	100%	99.6%	97.2%	{Hypogonadotropic hypogonadism 15 with or without anosmia}, 614880
IGSF10	100%	100%	100%	100%	99.7%	
IL17RD	100%	100%	100%	100%	99.3%	Hypogonadotropic hypogonadism 18 with or without anosmia, 615267
KISS1	100%	100%	100%	99.9%	98.5%	?Hypogonadotropic hypogonadism 13 with or without anosmia, 614842

KISS1R	100%	100%	100%	100%	99.5%	Hypogonadotropic hypogonadism 8 with or without anosmia, 614837;?Precocious puberty, central, 1, 176400
KLB	100%	100%	100%	100%	99.7%	
LEP	100%	100%	100%	99.7%	98.3%	Obesity, morbid, due to leptin deficiency, 614962
LEPR	94.6%	94.6%	100%	100%	99.8%	Obesity, morbid, due to leptin receptor deficiency, 614963
LHB	100%	100%	100%	100%	99.4%	Hypogonadotropic hypogonadism 23 with or without anosmia, 228300
LHX3	100%	100%	100%	100%	99.1%	Pituitary hormone deficiency, combined, 3, 221750
NDNF	100%	100%	100%	100%	99.9%	Hypogonadotropic hypogonadism 25 with anosmia, 618841
NOS1	100%	100%	100%	99.9%	98.6%	
NR0B1	100%	99.6%	98.7%	89%	71.8%	Adrenal hypoplasia, congenital, 300200;46XY sex reversal 2, dosage-sensitive, 300018

NSMF	100%	100%	100%	99.8%	98.8%	Hypogonadotropic hypogonadism 9 with or without anosmia, 614838
NTN1	100%	100%	100%	99.9%	98.7%	Mirror movements 4, 618264
PCSK1	100%	100%	100%	99.9%	99.5%	{Obesity, susceptibility to, BMIQ12}, 612362;Endocrinopathy due to proprotein convertase 1/3 deficiency, 600955
PHF6	100%	100%	99.6%	92.7%	76.3%	Borjeson-Forssman-Lehmann syndrome, 301900
PLXNA1	100%	100%	100%	100%	99.4%	Dworschak-Punetha neurodevelopmental syndrome, 619955
PNPLA6	100%	100%	100%	99.9%	98.8%	Spastic paraplegia 39, autosomal recessive, 612020;Oliver-McFarlane syndrome, 275400;?Laurence-Moon syndrome, 245800;Boucher-Neuhauser syndrome, 215470

POLG	100%	100%	100%	100%	99.5%	Mitochondrial recessive ataxia syndrome (includes SANDO and SCAE), 607459;Mitochondrial DNA depletion syndrome 4B (MNGIE type), 613662;Mitochondrial DNA depletion syndrome 4A (Alpers type), 203700;Progressive external ophthalmoplegia, autosomal dominant 1, 157640;Progressive external ophthalmoplegia, autosomal recessive 1, 258450
POLR3A	100%	100%	100%	100%	99.3%	Wiedemann-Rautenstrauch syndrome, 264090;Leukodystrophy, hypomyelinating, 7, with or without oligodontia and/or hypogonadotropic hypogonadism, 607694

POLR3B	100%	100%	100%	100%	99.8%	Leukodystrophy, hypomyelinating, 8, with or without oligodontia and/or hypogonadotropic hypogonadism, 614381;Charcot-Marie-Tooth disease, demyelinating, type 1I, 619742
POLR3GL	100%	100%	100%	100%	99.4%	Short stature, oligodontia, dysmorphic facies, and motor delay, 619234
PROK2	100%	100%	100%	100%	99.7%	Hypogonadotropic hypogonadism 4 with or without anosmia, 610628
PROKR2	100%	100%	100%	99.8%	99.3%	Hypogonadotropic hypogonadism 3 with or without anosmia, 244200
PROP1	100%	100%	100%	99.9%	98.8%	Pituitary hormone deficiency, combined, 2, 262600
SEMA3A	100%	100%	100%	100%	99.8%	{Hypogonadotropic hypogonadism 16 with or without anosmia}, 614897
SEMA3E	100%	100%	100%	100%	99.7%	

SOX10	97.8%	97.8%	100%	99.9%	98.3%	Waardenburg syndrome, type 4C, 613266;PCWH syndrome, 609136;Waardenburg syndrome, type 2E, with or without neurologic involvement, 611584
SOX11	100%	100%	100%	99.5%	94.3%	Intellectual developmental disorder with microcephaly and with or without ocular malformations or hypogonadotropic hypogonadism, 615866
SOX2	100%	99.9%	100%	99.9%	96.8%	Optic nerve hypoplasia and abnormalities of the central nervous system, 206900;Microphthalmia , syndromic 3, 206900
SPRY4	100%	100%	100%	100%	98.9%	Hypogonadotropic hypogonadism 17 with or without anosmia, 615266
TAC3	100%	100%	100%	99.7%	98.1%	Hypogonadotropic hypogonadism 10 with or without anosmia, 614839

TACR3	100%	100%	100%	100%	99.6%	Hypogonadotropic hypogonadism 11 with or without anosmia, 614840
TCF12	100%	100%	100%	100%	99.8%	Craniosynostosis 3, 615314;Hypogonadotropic hypogonadism 26 with or without anosmia, 619718
TENM1	100%	99.7%	99%	90.8%	72.8%	
WDR11	100%	100%	100%	100%	99.7%	Intellectual developmental disorder, autosomal recessive 78, 620237;Hypogonadotropic hypogonadism 14 with or without anosmia, 614858

Gene symbols used follow HGCN guidelines: Gray KA, Yates B, Seal RL, Wright MW, Bruford EA. Nucleic Acids Res. 2015 Jan 43(Database issue):D1079-85.

TWIST X2 covered 10x describes the percentage of a gene's coding sequence that is covered at least 10x when analyzed by WES using TWIST X2 chemistry mapped against GRCh38.

TWIST X2 covered 20x describes the percentage of a gene's coding sequence that is covered at least 20x when analyzed by WES using TWIST X2 chemistry mapped against GRCh38.

srWGS covered 10x describes the percentage of a gene's coding sequence that is covered at least 10x when analyzed by WGS mapped against GRCh38.

srWGS covered 15x describes the percentage of a gene's coding sequence that is covered at least 15x when analyzed by WGS mapped against GRCh38.

srWGS covered 20x describes the percentage of a gene's coding sequence that is covered at least 20x when analyzed by WGS mapped against GRCh38.

non-protein coding genes are covered, but as coverage statistics are based on protein coding regions, statistics could not be generated.

OMIM release used for OMIM disease identifiers and descriptions : November 25th, 2024.

This list is accurate for panel version DG 4.1.0

Ad 1. Blank field signifies a gene without a current OMIM association Ad 2. OMIM phenotype descriptions between {} signify risk factors