

PANEL HEREDITARY RENALCANCER DG-4.2.0 (12 GENES)

Gene	Twist X2 covered 10x	Twist X2 covered 20x	srWGS covered 10x	srWGS covered 15x	srWGS covered 20x	Associated Phenotype description and OMIM disease ID
BAP1	100%	100%	100%	100%	99%	Kury-Isidor syndrome, 619762;Tumor predisposition syndrome 1, 614327;{Uveal melanoma, susceptibility to, 2}, 606661
FH	100%	100%	100%	100%	99.3%	Leiomyomatosis and renal cell cancer, 150800;Fumarase deficiency, 606812
FLCN	100%	100%	100%	99.9%	98.9%	Birt-Hogg-Dube syndrome, 135150;Colorectal cancer, somatic, 114500;Pneumothorax, primary spontaneous, 173600;Renal carcinoma, chromophobe, somatic, 144700

MET	100%	100%	100%	99.9%	99.5%	Renal cell carcinoma, papillary, 1, familial and somatic, 605074;?Arthrogryposis , distal, type 11, 620019;Hepatocellular carcinoma, childhood type, somatic, 114550;{Osteofibrous dysplasia, susceptibility to}, 607278;?Deafness, autosomal recessive 97, 616705
PRDM10	100%	100%	100%	100%	99.2%	?Birt-Hogg-Dube syndrome 2, 620459
PTEN	94.5%	94.5%	100%	99.9%	99.2%	{Glioma susceptibility 2}, 613028;{Meningioma}, 607174;Cowden syndrome 1, 158350;Lhermitte-Duclos disease, 158350;Prostate cancer, somatic, 176807;Macrocephaly/autism syndrome, 605309

SDHA	100%	100%	100%	99.8%	99%	Cardiomyopathy, dilated, 1GG, 613642;Mitochondrial complex II deficiency, nuclear type 1, 252011;Neurodegeneration with ataxia and late-onset optic atrophy, 619259;Pheochromocytoma/paranglioma syndrome 5, 614165
SDHAF2	100%	99.2%	100%	100%	99.5%	Pheochromocytoma/paranglioma syndrome 2, 601650
SDHB	100%	100%	100%	99.8%	98.9%	Pheochromocytoma/paranglioma syndrome 4, 115310;Mitochondrial complex II deficiency, nuclear type 4, 619224;Gastrointestinal stromal tumor, 606764;Paranglioma and gastric stromal sarcoma, 606864
SDHC	100%	100%	100%	100%	99.2%	Pheochromocytoma/paranglioma syndrome 3, 605373;Paranglioma and gastric stromal sarcoma, 606864;Gastrointestinal stromal tumor, 606764

SDHD	78.9%	78.9%	100%	99.9%	98.1%	Pheochromocytoma/paraganglioma syndrome 1, 168000;Paraganglioma and gastric stromal sarcoma, 606864;Mitochondrial complex II deficiency, nuclear type 3, 619167
VHL	88%	87.7%	100%	99.9%	98.3%	Hemangioblastoma, cerebellar, somatic;Erythrocytosis, familial, 2, 263400;von Hippel-Lindau syndrome, 193300;Renal cell carcinoma, somatic, 144700;Pheochromocytoma, 171300

Gene symbols used follow HGCN guidelines: Gray KA, Yates B, Seal RL, Wright MW, Bruford EA. Nucleic Acids Res. 2015 Jan 43(Database issue):D1079-85.

TWIST X2 covered 10x describes the percentage of a gene's coding sequence that is covered at least 10x when analyzed by WES using TWIST X2 chemistry mapped against GRCh38.

TWIST X2 covered 20x describes the percentage of a gene's coding sequence that is covered at least 20x when analyzed by WES using TWIST X2 chemistry mapped against GRCh38.

srWGS covered 10x describes the percentage of a gene's coding sequence that is covered at least 10x when analyzed by WGS mapped against GRCh38.

srWGS covered 15x describes the percentage of a gene's coding sequence that is covered at least 15x when analyzed by WGS mapped against GRCh38.

srWGS covered 20x describes the percentage of a gene's coding sequence that is covered at least 20x when analyzed by WGS mapped against GRCh38.

non-protein coding genes are covered, but as coverage statistics are based on protein coding regions, statistics could not be generated.

OMIM release used for OMIM disease identifiers and descriptions : November 25th, 2024.

This list is accurate for panel version DG 4.2.0

Ad 1. Blank field signifies a gene without a current OMIM association Ad 2. OMIM phenotype descriptions between {} signify risk factors