

# MOVEMENT DISORDERS PANEL DG-4.1.0 (431 GENES)

| <i>Gene</i> | <i>Twist X2 covered &gt;10x</i> | <i>Twist X2 covered &gt;20x</i> | <i>WGS covered &gt;10x</i> | <i>WGS covered &gt;20x</i> | <i>Associated Phenotype description and OMIM disease ID</i>                                                            |
|-------------|---------------------------------|---------------------------------|----------------------------|----------------------------|------------------------------------------------------------------------------------------------------------------------|
| AARS2       | 100%                            | 100%                            | 100%                       | 99.3%                      | Leukoencephalopathy, progressive, with ovarian failure, 615889;Combined oxidative phosphorylation deficiency 8, 614096 |
| ABCB7       | 100%                            | 99.7%                           | 99.1%                      | 74.9%                      | Anemia, sideroblastic, with ataxia, 301310                                                                             |
| ABCD1       | 100%                            | 98.5%                           | 98.7%                      | 69.5%                      | Adrenoleukodystrophy, 300100;Adrenomyeloneuropathy, adult, 300100                                                      |
| ABHD12      | 100%                            | 100%                            | 100%                       | 99.6%                      | Polyneuropathy, hearing loss, ataxia, retinitis pigmentosa, and cataract, 612674                                       |
| ABHD16A     | 100%                            | 100%                            | 100%                       | 99%                        | Spastic paraplegia 86, autosomal recessive, 619735                                                                     |
| ACO2        | 93.1%                           | 90.3%                           | 100%                       | 99.3%                      | Optic atrophy 9, 616289;Infantile cerebellar-retinal degeneration, 614559                                              |

|       |       |       |      |       |                                                                                                                                                                                                                                                                                                              |
|-------|-------|-------|------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ACTB  | 100%  | 100%  | 100% | 99.3% | Baraitser-Winter syndrome 1, 243310;Becker nevus, syndromic or isolated, somatic mosaic, 604919;Thrombocytopenia 8, with dysmorphic features and developmental delay, 620475;Dystonia-deafness syndrome 1, 607371;Congenital smooth muscle hamartoma with or without hemihypertrophy, somatic mosaic, 620470 |
| ADAR  | 100%  | 100%  | 100% | 99.2% | Dyschromatosis symmetrica hereditaria, 127400;Aicardi-Goutieres syndrome 6, 615010                                                                                                                                                                                                                           |
| ADCY5 | 97.4% | 97.1% | 100% | 98.2% | Dyskinesia with orofacial involvement, autosomal dominant, 606703;Neurodevelopmental disorder with hyperkinetic movements and dyskinesia, 619651;Dyskinesia with orofacial involvement, autosomal recessive, 619647                                                                                          |

|         |      |      |      |       |                                                                                                                                                                                                      |
|---------|------|------|------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ADGRG1  | 100% | 100% | 100% | 98.8% | Cortical dysplasia, complex, with other brain malformations 14B, (bilateral perisylvian), 615752;Cortical dysplasia, complex, with other brain malformations 14A, (bilateral frontoparietal), 606854 |
| ADPRS   | 100% | 100% | 100% | 98.6% | Neurodegeneration, childhood-onset, stress-induced, with variable ataxia and seizures, 618170                                                                                                        |
| AFG3L2  | 100% | 100% | 100% | 99.4% | Spastic ataxia 5, autosomal recessive, 614487;Optic atrophy 12, 618977;Spinocerebellar ataxia 28, 610246                                                                                             |
| AGA     | 100% | 100% | 100% | 99.4% | Aspartylglucosaminuria, 208400                                                                                                                                                                       |
| AGTPBP1 | 100% | 100% | 100% | 99.7% | Neurodegeneration, childhood-onset, with cerebellar atrophy, 618276                                                                                                                                  |
| AIMP1   | 100% | 100% | 100% | 99.6% | Leukodystrophy, hypomyelinating, 3, 260600                                                                                                                                                           |

|          |       |       |      |       |                                                                                                                                                                                                  |
|----------|-------|-------|------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ALDH18A1 | 100%  | 100%  | 100% | 99.6% | Spastic paraplegia 9A, autosomal dominant, 601162;Cutis laxa, autosomal recessive, type IIIA, 219150;Spastic paraplegia 9B, autosomal recessive, 616586;Cutis laxa, autosomal dominant 3, 616603 |
| ALDH3A2  | 93.5% | 93.5% | 100% | 99.6% | Sjogren-Larsson syndrome, 270200                                                                                                                                                                 |
| ALDH5A1  | 100%  | 100%  | 100% | 98.9% | Succinic semialdehyde dehydrogenase deficiency, 271980                                                                                                                                           |
| ALS2     | 97.1% | 97%   | 100% | 99.6% | Primary lateral sclerosis, juvenile, 606353;Spastic paralysis, infantile onset ascending, 607225;Amyotrophic lateral sclerosis 2, juvenile, 205100                                               |
| AMFR     | 100%  | 100%  | 100% | 98.4% | Spastic paraplegia 89, autosomal recessive, 620379                                                                                                                                               |
| AMPD2    | 100%  | 100%  | 100% | 99.1% | Pontocerebellar hypoplasia, type 9, 615809;?Spastic paraplegia 63, autosomal recessive, 615686                                                                                                   |
| ANO10    | 100%  | 100%  | 100% | 99.7% | Spinocerebellar ataxia, autosomal recessive 10, 613728                                                                                                                                           |
| ANO3     | 100%  | 100%  | 100% | 99.7% | Dystonia 24, 615034                                                                                                                                                                              |
| AOPEP    | 100%  | 100%  | 100% | 99.3% | Dystonia 31, 619565                                                                                                                                                                              |

|       |       |       |      |       |                                                                                               |
|-------|-------|-------|------|-------|-----------------------------------------------------------------------------------------------|
| AP4B1 | 100%  | 100%  | 100% | 99.5% | Spastic paraplegia 47, autosomal recessive, 614066                                            |
| AP4E1 | 100%  | 100%  | 100% | 99.7% | Stuttering, familial persistent, 1, 184450;Spastic paraplegia 51, autosomal recessive, 613744 |
| AP4M1 | 100%  | 100%  | 100% | 98.7% | Spastic paraplegia 50, autosomal recessive, 612936                                            |
| AP4S1 | 87.8% | 87.5% | 100% | 99.6% | Spastic paraplegia 52, autosomal recessive, 614067                                            |
| AP5Z1 | 100%  | 100%  | 100% | 98.7% | Spastic paraplegia 48, autosomal recessive, 613647                                            |
| APTX  | 100%  | 100%  | 100% | 99.3% | Ataxia, early-onset, with oculomotor apraxia and hypoalbuminemia, 208920                      |
| ARG1  | 93%   | 93%   | 100% | 99.8% | Argininemia, 207800                                                                           |
| ARSA  | 100%  | 100%  | 100% | 98.7% | Metachromatic leukodystrophy, 250100                                                          |

|       |       |       |       |       |                                                                                                                                                                                                                                                    |
|-------|-------|-------|-------|-------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ARX   | 98.5% | 94.3% | 91.9% | 52.1% | Proud syndrome, 300004;Hydranencephaly with abnormal genitalia, 300215;Partington syndrome, 309510;Developmental and epileptic encephalopathy 1, 308350;Lissencephaly, X-linked 2, 300215;Intellectual developmental disorder, X-linked 29, 300419 |
| ASPA  | 100%  | 100%  | 100%  | 99.8% | Canavan disease, 271900                                                                                                                                                                                                                            |
| ATCAY | 100%  | 100%  | 100%  | 98.9% | Ataxia, cerebellar, Cayman type, 601238                                                                                                                                                                                                            |
| ATG7  | 100%  | 100%  | 100%  | 99.8% | Spinocerebellar ataxia, autosomal recessive 31, 619422                                                                                                                                                                                             |
| ATL1  | 100%  | 100%  | 100%  | 99.7% | Spastic paraplegia 3A, autosomal dominant, 182600;Neuropathy, hereditary sensory, type ID, 613708                                                                                                                                                  |
| ATM   | 100%  | 100%  | 100%  | 99.5% | Lymphoma, B-cell non-Hodgkin, somatic;Ataxia-telangiectasia, 208900;{Breast cancer, susceptibility to}, 114480;T-cell prolymphocytic leukemia, somatic;Lymphoma, mantle cell, somatic                                                              |

|         |      |       |       |       |                                                                                                                                                                                                                                                                                        |
|---------|------|-------|-------|-------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ATN1    | 100% | 100%  | 99.9% | 95.7% | Dentatorubral-pallidoluysian atrophy, 125370;Congenital hypotonia, epilepsy, developmental delay, and digital anomalies, 618494                                                                                                                                                        |
| ATP13A2 | 100% | 100%  | 100%  | 99.2% | Spastic paraplegia 78, autosomal recessive, 617225;Kufor-Rakeb syndrome, 606693                                                                                                                                                                                                        |
| ATP1A2  | 100% | 100%  | 100%  | 99.1% | Developmental and epileptic encephalopathy 98, 619605;Fetal akinesia, respiratory insufficiency, microcephaly, polymicrogyria, and dysmorphic facies, 619602;Alternating hemiplegia of childhood 1, 104290;Migraine, familial basilar, 602481;Migraine, familial hemiplegic, 2, 602481 |
| ATP1A3  | 100% | 100%  | 100%  | 98.5% | Alternating hemiplegia of childhood 2, 614820;Dystonia-12, 128235;CAPOS syndrome, 601338;Developmental and epileptic encephalopathy 99, 619606                                                                                                                                         |
| ATP2B3  | 100% | 99.3% | 97.9% | 67.7% | ?Spinocerebellar ataxia, X-linked 1, 302500                                                                                                                                                                                                                                            |
| ATP7B   | 100% | 100%  | 100%  | 99.7% | Wilson disease, 277900                                                                                                                                                                                                                                                                 |

|          |       |       |       |       |                                                                                                                                                            |
|----------|-------|-------|-------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| B4GALNT1 | 100%  | 100%  | 100%  | 99%   | Spastic paraplegia 26, autosomal recessive, 609195                                                                                                         |
| BCAP31   | 98.9% | 86.5% | 98.5% | 69.8% | Deafness, dystonia, and cerebral hypomyelination, 300475                                                                                                   |
| BCKDHA   | 100%  | 100%  | 100%  | 98.4% | Maple syrup urine disease, type Ia, 248600                                                                                                                 |
| BCKDHB   | 100%  | 100%  | 100%  | 99.2% | Maple syrup urine disease, type Ib, 620698                                                                                                                 |
| BCL11B   | 99.9% | 99.3% | 100%  | 94.8% | Immunodeficiency 49, severe combined, 617237;Intellectual developmental disorder with dysmorphic facies, speech delay, and T-cell abnormalities, 618092    |
| BRAT1    | 100%  | 100%  | 100%  | 99.2% | Neurodevelopmental disorder with cerebellar atrophy and with or without seizures, 618056;Rigidity and multifocal seizure syndrome, lethal neonatal, 614498 |



|          |       |       |      |       |                                                                                                                                                                                                                                              |
|----------|-------|-------|------|-------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| BSCL2    | 100%  | 100%  | 100% | 98.9% | Lipodystrophy, congenital generalized, type 2, 269700;Neuronopathy, distal hereditary motor, autosomal dominant 13, 619112;Silver spastic paraplegia syndrome, 270685;Encephalopathy, progressive, with or without lipodystrophy, 615924     |
| BTB      | 94.2% | 94.2% | 100% | 99.7% | Biotinidase deficiency, 253260                                                                                                                                                                                                               |
| C19orf12 | 100%  | 100%  | 100% | 98.5% | Neurodegeneration with brain iron accumulation 4, 614298;?Spastic paraplegia 43, autosomal recessive, 615043                                                                                                                                 |
| CA8      | 100%  | 100%  | 100% | 99.8% | Spinocerebellar ataxia, autosomal recessive 34, 613227                                                                                                                                                                                       |
| CACNA1A  | 100%  | 100%  | 100% | 98%   | Spinocerebellar ataxia 6, 183086;Episodic ataxia, type 2, 108500;Developmental and epileptic encephalopathy 42, 617106;Migraine, familial hemiplegic, 1, with progressive cerebellar ataxia, 141500;Migraine, familial hemiplegic, 1, 141500 |
| CACNA1E  | 100%  | 100%  | 100% | 99.2% | Developmental and epileptic encephalopathy 69, 618285                                                                                                                                                                                        |

|         |       |       |      |       |                                                                                                                                                                |
|---------|-------|-------|------|-------|----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CACNA1G | 100%  | 100%  | 100% | 98.8% | Spinocerebellar ataxia 42, 616795;Spinocerebellar ataxia 42, early-onset, severe, with neurodevelopmental deficits, 618087                                     |
| CACNB4  | 100%  | 100%  | 100% | 99.6% | {Epilepsy, juvenile myoclonic, susceptibility to, 6}, 607682;?Episodic ataxia, type 5, 613855;{Epilepsy, idiopathic generalized, susceptibility to, 9}, 607682 |
| CAMTA1  | 100%  | 100%  | 100% | 99.2% | Cerebellar dysfunction with variable cognitive and behavioral abnormalities, 614756                                                                            |
| CAPN1   | 100%  | 100%  | 100% | 99.1% | Spastic paraplegia 76, autosomal recessive, 616907                                                                                                             |
| CC2D2A  | 98.2% | 98.2% | 100% | 99.6% | COACH syndrome 2, 619111;Retinitis pigmentosa 93, 619845;Meckel syndrome 6, 612284;Joubert syndrome 9, 612285                                                  |
| CCT5    | 100%  | 100%  | 100% | 99.5% | ?Neuropathy, hereditary sensory, with spastic paraplegia, 256840                                                                                               |
| CHMP1A  | 100%  | 100%  | 100% | 98.8% | Pontocerebellar hypoplasia, type 8, 614961                                                                                                                     |

|       |      |       |       |       |                                                                                                                                                                                                                                                                                  |
|-------|------|-------|-------|-------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CLCN2 | 100% | 100%  | 100%  | 99.4% | Leukoencephalopathy with ataxia, 615651;Hyperaldosteronism , familial, type II, 605635;{Epilepsy, juvenile myoclonic, susceptibility to, 8}, 607628;{Epilepsy, juvenile absence, susceptibility to, 2}, 607628;{Epilepsy, idiopathic generalized, susceptibility to, 11}, 607628 |
| CLCN4 | 100% | 99.6% | 98.8% | 69%   | Raynaud-Claes syndrome, 300114                                                                                                                                                                                                                                                   |
| CLCN6 | 100% | 100%  | 100%  | 99.1% | Ceroid lipofuscinosis, neuronal, 15, 619173                                                                                                                                                                                                                                      |
| CLN5  | 83%  | 83%   | 100%  | 99.7% | Ceroid lipofuscinosis, neuronal, 5, 256731                                                                                                                                                                                                                                       |
| CLN6  | 100% | 100%  | 100%  | 98.8% | Ceroid lipofuscinosis, neuronal, 6B (Kufs type), 204300;Ceroid lipofuscinosis, neuronal, 6A, 601780                                                                                                                                                                              |
| CLP1  | 100% | 100%  | 100%  | 98.8% | Pontocerebellar hypoplasia, type 10, 615803                                                                                                                                                                                                                                      |

|        |      |      |      |       |                                                                                                                                                                                                                                                                                                                                 |
|--------|------|------|------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CLPB   | 100% | 100% | 100% | 98.9% | Neutropenia, severe congenital, 9, autosomal dominant, 619813;3-methylglutaconic aciduria, type VIIB, autosomal recessive, 616271;3-methylglutaconic aciduria, type VIIA, autosomal dominant, 619835                                                                                                                            |
| COASY  | 100% | 100% | 100% | 98.9% | Pontocerebellar hypoplasia, type 12, 618266;Neurodegeneration with brain iron accumulation 6, 615643                                                                                                                                                                                                                            |
| COL4A1 | 100% | 100% | 100% | 99.3% | ?Retinal arteries, tortuosity of, 180000;{Hemorrhage, intracerebral, susceptibility to}, 614519;Angiopathy, hereditary, with nephropathy, aneurysms, and muscle cramps, 611773;Microangiopathy and leukoencephalopathy, pontine, autosomal dominant, 618564;Brain small vessel disease with or without ocular anomalies, 175780 |
| COL4A2 | 100% | 100% | 100% | 98.7% | Brain small vessel disease 2, 614483;{Hemorrhage, intracerebral, susceptibility to}, 614519                                                                                                                                                                                                                                     |

|        |       |       |      |       |                                                                                                                |
|--------|-------|-------|------|-------|----------------------------------------------------------------------------------------------------------------|
| COL6A1 | 100%  | 100%  | 100% | 98.7% | Ullrich congenital muscular dystrophy 1A, 254090;Bethlem myopathy 1A, 158810                                   |
| COL6A2 | 100%  | 100%  | 100% | 99%   | ?Myosclerosis, congenital, 255600;Ullrich congenital muscular dystrophy 1B, 620727;Bethlem myopathy 1B, 620725 |
| COL6A3 | 100%  | 100%  | 100% | 99.6% | Bethlem myopathy 1C, 620726;Ullrich congenital muscular dystrophy 1C, 620728;Dystonia 27, 616411               |
| COQ2   | 96.3% | 96.3% | 100% | 98.7% | {Multiple system atrophy, susceptibility to}, 146500;Coenzyme Q10 deficiency, primary, 1, 607426               |
| COQ4   | 100%  | 100%  | 100% | 98.7% | Coenzyme Q10 deficiency, primary, 7, 616276;Spastic ataxia 10, autosomal recessive, 620666                     |
| COQ8A  | 100%  | 100%  | 100% | 98.7% | Coenzyme Q10 deficiency, primary, 4, 612016                                                                    |
| COQ9   | 100%  | 100%  | 100% | 98.6% | Coenzyme Q10 deficiency, primary, 5, 614654                                                                    |
| COX20  | 100%  | 100%  | 100% | 99.6% | Mitochondrial complex IV deficiency, nuclear type 11, 619054                                                   |
| CP     | 100%  | 100%  | 100% | 99.2% | Aceruloplasminemia, 604290                                                                                     |

|         |       |       |       |       |                                                                                                                                         |
|---------|-------|-------|-------|-------|-----------------------------------------------------------------------------------------------------------------------------------------|
| CSF1R   | 100%  | 100%  | 100%  | 99.2% | Brain abnormalities, neurodegeneration, and dysosteosclerosis, 618476;Leukoencephalopathy, diffuse hereditary, with spheroids 1, 221820 |
| CSTB    | 100%  | 100%  | 100%  | 99.5% | Epilepsy, progressive myoclonic 1A (Unverricht and Lundborg), 254800                                                                    |
| CTBP1   | 99.3% | 97.9% | 99.9% | 97.1% | Hypotonia, ataxia, developmental delay, and tooth enamel defect syndrome, 617915                                                        |
| CTC1    | 100%  | 100%  | 100%  | 99.3% | Cerebroretinal microangiopathy with calcifications and cysts, 612199                                                                    |
| CTSD    | 100%  | 100%  | 100%  | 99.4% | Ceroid lipofuscinosis, neuronal, 10, 610127                                                                                             |
| CTSF    | 100%  | 100%  | 100%  | 99.1% | Ceroid lipofuscinosis, neuronal, 13 (Kufs type), 615362                                                                                 |
| CWF19L1 | 100%  | 100%  | 100%  | 99.7% | Spinocerebellar ataxia, autosomal recessive 17, 616127                                                                                  |
| CYP27A1 | 100%  | 100%  | 100%  | 99%   | Cerebrotendinous xanthomatosis, 213700                                                                                                  |
| CYP2U1  | 100%  | 100%  | 100%  | 97%   | Spastic paraplegia 56, autosomal recessive, 615030                                                                                      |

|        |      |      |      |       |                                                                                                                                                                                                                   |
|--------|------|------|------|-------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CYP7B1 | 100% | 100% | 100% | 99.7% | Spastic paraplegia 5A, autosomal recessive, 270800;Bile acid synthesis defect, congenital, 3, 613812                                                                                                              |
| DARS1  | 100% | 100% | 100% | 99.9% | Hypomyelination with brainstem and spinal cord involvement and leg spasticity, 615281                                                                                                                             |
| DARS2  | 100% | 100% | 100% | 99.5% | Leukoencephalopathy with brain stem and spinal cord involvement and lactate elevation, 611105                                                                                                                     |
| DBT    | 100% | 100% | 100% | 99.5% | Maple syrup urine disease, type II, 620699                                                                                                                                                                        |
| DCAF17 | 100% | 100% | 100% | 99.4% | Woodhouse-Sakati syndrome, 241080                                                                                                                                                                                 |
| DCC    | 100% | 100% | 100% | 99.2% | Mirror movements 1 and/or agenesis of the corpus callosum, 157600;Esophageal carcinoma, somatic, 133239;Colorectal cancer, somatic, 114500;Gaze palsy, familial horizontal, with progressive scoliosis, 2, 617542 |

|       |       |       |      |       |                                                                                                                                                                         |
|-------|-------|-------|------|-------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| DCTN1 | 100%  | 100%  | 100% | 98.9% | Perry syndrome, 168605;{Amyotrophic lateral sclerosis, susceptibility to}, 105400;Neuronopathy, distal hereditary motor, autosomal dominant 14, 607641                  |
| DDC   | 100%  | 100%  | 100% | 99.2% | Aromatic L-amino acid decarboxylase deficiency, 608643                                                                                                                  |
| DDHD1 | 100%  | 100%  | 100% | 98.7% | Spastic paraplegia 28, autosomal recessive, 609340                                                                                                                      |
| DDHD2 | 100%  | 100%  | 100% | 99.8% | Spastic paraplegia 54, autosomal recessive, 615033                                                                                                                      |
| DEGS1 | 100%  | 100%  | 100% | 99.6% | Leukodystrophy, hypomyelinating, 18, 618404                                                                                                                             |
| DHDDS | 73.7% | 73.7% | 100% | 99%   | Developmental delay and seizures with or without movement abnormalities, 617836;?Congenital disorder of glycosylation, type 1bb, 613861;Retinitis pigmentosa 59, 613861 |
| DLAT  | 100%  | 100%  | 100% | 99.7% | Pyruvate dehydrogenase E2 deficiency, 245348                                                                                                                            |
| DLD   | 100%  | 100%  | 100% | 99.7% | Dihydrolipoamide dehydrogenase deficiency, 246900                                                                                                                       |



|         |       |       |      |       |                                                                                                                         |
|---------|-------|-------|------|-------|-------------------------------------------------------------------------------------------------------------------------|
| DNAJC12 | 100%  | 100%  | 100% | 99.6% | Hyperphenylalaninemia, mild, non-BH4-deficient, 617384                                                                  |
| DNAJC3  | 100%  | 100%  | 100% | 99.9% | Ataxia, combined cerebellar and peripheral, with hearing loss and diabetes mellitus, 616192                             |
| DNAL4   | 100%  | 100%  | 100% | 97.5% | ?Mirror movements 3, 616059                                                                                             |
| DNM1L   | 100%  | 100%  | 100% | 99.6% | Optic atrophy 5, 610708;Encephalopathy, lethal, due to defective mitochondrial peroxisomal fission 1, 614388            |
| DNMT1   | 99.9% | 99.1% | 100% | 99.4% | Neuropathy, hereditary sensory, type IE, 614116;Cerebellar ataxia, deafness, and narcolepsy, autosomal dominant, 604121 |
| DPYS    | 100%  | 100%  | 100% | 99.4% | Dihydropyrimidinuria, 222748                                                                                            |
| DSTYK   | 100%  | 100%  | 100% | 99.3% | Spastic paraplegia 23, autosomal recessive, 270750;Congenital anomalies of kidney and urinary tract 1, 610805           |
| DTYMK   | 100%  | 100%  | 100% | 99.1% | Neurodegeneration, childhood-onset, with progressive microcephaly, 619847                                               |

|         |      |      |      |       |                                                                                                                   |
|---------|------|------|------|-------|-------------------------------------------------------------------------------------------------------------------|
| EBF3    | 100% | 100% | 100% | 98.1% | Hypotonia, ataxia, and delayed development syndrome, 617330                                                       |
| ECHS1   | 100% | 100% | 100% | 99.4% | Mitochondrial short-chain enoyl-CoA hydratase 1 deficiency, 616277                                                |
| EIF2AK1 | 100% | 100% | 100% | 99.7% | ?Leukoencephalopathy, motor delay, spasticity, and dysarthria syndrome, 618878                                    |
| EIF2AK2 | 100% | 100% | 100% | 99.6% | Leukoencephalopathy, developmental delay, and episodic neurologic regression syndrome, 618877;Dystonia 33, 619687 |
| EIF2B1  | 100% | 100% | 100% | 99.3% | Leukoencephalopathy with vanishing white matter 1, with or without ovarian failure, 603896                        |
| EIF2B2  | 100% | 100% | 100% | 97.8% | Leukoencephalopathy with vanishing white matter 2, with or without ovarian failure, 620312                        |
| EIF2B3  | 100% | 100% | 100% | 99.3% | Leukoencephalopathy with vanishing white matter 3, with or without ovarian failure, 620313                        |
| EIF2B4  | 100% | 100% | 100% | 99.6% | Leukoencephalopathy with vanishing white matter 4, with or without ovarian failure, 620314                        |

|        |       |       |      |       |                                                                                                                                               |
|--------|-------|-------|------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------|
| EIF2B5 | 100%  | 100%  | 100% | 99.4% | Leukoencephalopathy with vanishing white matter 5, with or without ovarian failure, 620315                                                    |
| ELOVL1 | 100%  | 100%  | 100% | 99.1% | Ichthyotic keratoderma, spasticity, hypomyelination, and dysmorphic facies, 618527                                                            |
| ELOVL4 | 100%  | 100%  | 100% | 99%   | Spinocerebellar ataxia 34, 133190;Stargardt disease 3, 600110;Ichthyosis, spastic quadriplegia, and impaired intellectual development, 614457 |
| ELOVL5 | 100%  | 100%  | 100% | 99.4% | Spinocerebellar ataxia 38, 615957                                                                                                             |
| ENTPD1 | 100%  | 100%  | 100% | 99.7% | Spastic paraplegia 64, autosomal recessive, 615683                                                                                            |
| ERCC2  | 99.9% | 98.8% | 100% | 98.9% | Xeroderma pigmentosum, group D, 278730;Trichothiodystrophy 1, photosensitive, 601675;?Cerebrooculofacio skeletal syndrome 2, 610756           |

|        |      |       |      |       |                                                                                                                                                                               |
|--------|------|-------|------|-------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ERCC4  | 100% | 100%  | 100% | 99.5% | Xeroderma pigmentosum, type F/Cockayne syndrome, 278760;XFE progeroid syndrome, 610965;Xeroderma pigmentosum, group F, 278760;Fanconi anemia, complementation group Q, 615272 |
| ERLIN1 | 100% | 100%  | 100% | 99%   | Spastic paraplegia 62, autosomal recessive, 615681                                                                                                                            |
| ERLIN2 | 100% | 100%  | 100% | 99.2% | Spastic paraplegia 18A, autosomal dominant, 620512;Spastic paraplegia 18B, autosomal recessive, 611225                                                                        |
| ETHE1  | 100% | 100%  | 100% | 99.3% | Ethylmalonic encephalopathy, 602473                                                                                                                                           |
| EXOSC3 | 100% | 100%  | 100% | 99.6% | Pontocerebellar hypoplasia, type 1B, 614678                                                                                                                                   |
| EXOSC5 | 100% | 100%  | 100% | 97%   | Cerebellar ataxia, brain abnormalities, and cardiac conduction defects, 619576                                                                                                |
| EXOSC8 | 100% | 100%  | 100% | 99.9% | Pontocerebellar hypoplasia, type 1C, 616081                                                                                                                                   |
| EXOSC9 | 100% | 99.9% | 100% | 99.3% | Pontocerebellar hypoplasia, type 1D, 618065                                                                                                                                   |
| FA2H   | 100% | 100%  | 100% | 99.1% | Spastic paraplegia 35, autosomal recessive, 612319                                                                                                                            |

|        |      |      |      |       |                                                                                                                  |
|--------|------|------|------|-------|------------------------------------------------------------------------------------------------------------------|
| FAR1   | 100% | 100% | 100% | 99.9% | Peroxisomal fatty acyl-CoA reductase 1 disorder, 616154;Cataracts, spastic paraparesis, and speech delay, 619338 |
| FARS2  | 100% | 100% | 100% | 99%   | Combined oxidative phosphorylation deficiency 14, 614946;Spastic paraplegia 77, autosomal recessive, 617046      |
| FBXO7  | 100% | 100% | 100% | 99.6% | Parkinson disease 15, autosomal recessive, 260300                                                                |
| FGF14  | 100% | 100% | 100% | 99.3% | Spinocerebellar ataxia 27A, 193003;Spinocerebellar ataxia 27B, late-onset, 620174                                |
| FICD   | 100% | 100% | 100% | 99.4% | Spastic paraplegia 92, autosomal recessive, 620911                                                               |
| FLVCR1 | 100% | 100% | 100% | 99.2% | Ataxia, posterior column, with retinitis pigmentosa, 609033                                                      |
| FOLR1  | 100% | 100% | 100% | 98.9% | Neurodegeneration due to cerebral folate transport deficiency, 613068                                            |
| FRMD5  | 100% | 100% | 100% | 99.2% | Neurodevelopmental disorder with eye movement abnormalities and ataxia, 620094                                   |

|       |      |       |       |       |                                                                                                                                                            |
|-------|------|-------|-------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| FRMD7 | 100% | 99.8% | 98.6% | 73.9% | Nystagmus, infantile periodic alternating, X-linked, 310700;Nystagmus 1, congenital, X-linked, 310700                                                      |
| FTH1  | 100% | 100%  | 100%  | 98.5% | Neurodegeneration with brain iron accumulation 9, 620669;?Hemochromatosis, type 5, 615517                                                                  |
| FTL   | 100% | 99.9% | 100%  | 99%   | Hyperferritinemia-cataract syndrome, 600886;L-ferritin deficiency, dominant and recessive, 615604;Neurodegeneration with brain iron accumulation 3, 606159 |
| GALC  | 100% | 100%  | 100%  | 99.1% | Krabbe disease, 245200                                                                                                                                     |
| GAMT  | 100% | 100%  | 100%  | 98.2% | Cerebral creatine deficiency syndrome 2, 612736                                                                                                            |
| GAN   | 100% | 100%  | 100%  | 99.7% | Giant axonal neuropathy-1, 256850                                                                                                                          |

|       |      |      |      |       |                                                                                                                                                                                                                                                                                            |
|-------|------|------|------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| GBA1  | 100% | 100% | 100% | 99.1% | {Lewy body dementia, susceptibility to}, 127750;Gaucher disease, type II, 230900;Gaucher disease, type IIIC, 231005;Gaucher disease, type III, 231000;Gaucher disease, type I, 230800;Gaucher disease, perinatal lethal, 608013;{Parkinson disease, late-onset, susceptibility to}, 168600 |
| GBA2  | 100% | 100% | 100% | 99.3% | Spastic paraplegia 46, autosomal recessive, 614409                                                                                                                                                                                                                                         |
| GBE1  | 100% | 100% | 100% | 99.8% | Glycogen storage disease IV, 232500;Polyglucosan body disease, adult form, 263570                                                                                                                                                                                                          |
| GCDH  | 100% | 100% | 100% | 99.2% | Glutaricaciduria, type I, 231670                                                                                                                                                                                                                                                           |
| GCH1  | 100% | 100% | 100% | 99.4% | Dystonia, DOPA-responsive, 128230;Hyperphenylalanine mia, BH4-deficient, B, 233910                                                                                                                                                                                                         |
| GDAP2 | 100% | 100% | 100% | 99.6% | Spinocerebellar ataxia, autosomal recessive 27, 618369                                                                                                                                                                                                                                     |
| GFAP  | 100% | 100% | 100% | 99.6% | Alexander disease, 203450                                                                                                                                                                                                                                                                  |

|       |       |       |       |       |                                                                                                                                                                                                                                               |
|-------|-------|-------|-------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| GJB1  | 100%  | 100%  | 95.7% | 60.5% | Charcot-Marie-Tooth neuropathy, X-linked dominant, 1, 302800                                                                                                                                                                                  |
| GJC2  | 99.9% | 97.5% | 100%  | 97.2% | Lymphatic malformation 3, 613480;?Spastic paraplegia 44, autosomal recessive, 613206;Leukodystrophy, hypomyelinating, 2, 608804                                                                                                               |
| GLB1  | 100%  | 100%  | 100%  | 99.5% | GM1-gangliosidosis, type I, 230500;GM1-gangliosidosis, type III, 230650;Mucopolysaccharidosis type IVB (Morquio), 253010;GM1-gangliosidosis, type II, 230600                                                                                  |
| GLS   | 100%  | 100%  | 100%  | 99.5% | Global developmental delay, progressive ataxia, and elevated glutamine, 618412;?Infantile cataract, skin abnormalities, glutamate excess, and impaired intellectual development, 618339;Developmental and epileptic encephalopathy 71, 618328 |
| GNAL  | 100%  | 100%  | 100%  | 99.4% | Dystonia 25, 615073                                                                                                                                                                                                                           |
| GNAO1 | 100%  | 100%  | 100%  | 99.4% | Developmental and epileptic encephalopathy 17, 615473;Neurodevelopmental disorder with involuntary movements, 617493                                                                                                                          |



|        |      |       |       |       |                                                                                                                                                                                                                                                                                 |
|--------|------|-------|-------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| GOSR2  | 100% | 100%  | 100%  | 99.2% | Epilepsy, progressive myoclonic 6, 614018;Muscular dystrophy, congenital, with or without seizures, 620166                                                                                                                                                                      |
| GPR143 | 100% | 99.8% | 98.9% | 72.4% | Ocular albinism, type I, Nettleship-Falls type, 300500;Nystagmus 6, congenital, X-linked, 300814                                                                                                                                                                                |
| GRID2  | 100% | 100%  | 100%  | 99.5% | Spinocerebellar ataxia, autosomal recessive 18, 616204                                                                                                                                                                                                                          |
| GRIN1  | 100% | 100%  | 100%  | 98.7% | Neurodevelopmental disorder with or without hyperkinetic movements and seizures, autosomal recessive, 617820;Developmental and epileptic encephalopathy 101, 619814;Neurodevelopmental disorder with or without hyperkinetic movements and seizures, autosomal dominant, 614254 |
| GRIN2B | 100% | 100%  | 100%  | 99.4% | Developmental and epileptic encephalopathy 27, 616139;Intellectual developmental disorder, autosomal dominant 6, with or without seizures, 613970                                                                                                                               |

|       |      |      |      |       |                                                                                                                                                                                                                                                           |
|-------|------|------|------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| GRM1  | 100% | 100% | 100% | 99.4% | Spinocerebellar ataxia, autosomal recessive 13, 614831; Spinocerebellar ataxia 44, 617691                                                                                                                                                                 |
| GRN   | 100% | 100% | 100% | 99.8% | Frontotemporal dementia 2, 607485; Aphasia, primary progressive, 607485; Ceroid lipofuscinosis, neuronal, 11, 614706                                                                                                                                      |
| HACE1 | 100% | 100% | 100% | 99.3% | Spastic paraplegia and psychomotor retardation with or without seizures, 616756                                                                                                                                                                           |
| HEXB  | 100% | 100% | 100% | 98.8% | Sandhoff disease, infantile, juvenile, and adult forms, 268800                                                                                                                                                                                            |
| HK1   | 100% | 100% | 100% | 99.4% | Anemia, congenital, nonspherocytic hemolytic, 5, hexokinase deficient, 235700; Retinitis pigmentosa 79, 617460; Neuropathy, hereditary motor and sensory, Russe type, 605285; Neurodevelopmental disorder with visual defects and brain anomalies, 618547 |
| HPCA  | 100% | 100% | 100% | 98.3% | Dystonia 2, torsion, autosomal recessive, 224500                                                                                                                                                                                                          |

|         |       |       |       |       |                                                                                                                                                         |
|---------|-------|-------|-------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
| HPDL    | 100%  | 100%  | 100%  | 98.9% | Neurodevelopmental disorder with progressive spasticity and brain white matter abnormalities, 619026;Spastic paraplegia 83, autosomal recessive, 619027 |
| HPRT1   | 100%  | 99.8% | 99.1% | 73.6% | Hyperuricemia, HRPT-related, 300323;Lesch-Nyhan syndrome, 300322                                                                                        |
| HSD17B4 | 100%  | 100%  | 100%  | 99.6% | D-bifunctional protein deficiency, 261515;Perrault syndrome 1, 233400                                                                                   |
| HSPD1   | 99.6% | 97.6% | 100%  | 99.8% | Spastic paraplegia 13, autosomal dominant, 605280;Leukodystrophy, hypomyelinating, 4, 612233                                                            |
| HYCC1   | 100%  | 100%  | 100%  | 99.5% | Leukodystrophy, hypomyelinating, 5, 610532                                                                                                              |
| IBA57   | 100%  | 100%  | 100%  | 99.1% | Multiple mitochondrial dysfunctions syndrome 3, 615330;?Spastic paraplegia 74, autosomal recessive, 616451                                              |
| IRF2BPL | 100%  | 100%  | 98.2% | 89.6% | Neurodevelopmental disorder with regression, abnormal movements, loss of speech, and seizures, 618088                                                   |
| ISCA2   | 100%  | 100%  | 100%  | 99.5% | Multiple mitochondrial dysfunctions syndrome 4, 616370                                                                                                  |

|        |       |       |       |       |                                                                                                                           |
|--------|-------|-------|-------|-------|---------------------------------------------------------------------------------------------------------------------------|
| ITPR1  | 100%  | 100%  | 100%  | 99.5% | Gillespie syndrome, 206700;Spinocerebellar ataxia 29, congenital nonprogressive, 117360;Spinocerebellar ataxia 15, 606658 |
| JAM2   | 92.5% | 92%   | 100%  | 99.6% | Basal ganglia calcification, idiopathic, 8, autosomal recessive, 618824                                                   |
| JAM3   | 100%  | 100%  | 100%  | 99.8% | Hemorrhagic destruction of the brain, subependymal calcification, and cataracts, 613730                                   |
| KATNB1 | 100%  | 100%  | 100%  | 98.8% | Lissencephaly 6, with microcephaly, 616212                                                                                |
| KCNA1  | 100%  | 100%  | 100%  | 99.1% | Episodic ataxia/myokymia syndrome, 160120                                                                                 |
| KCNA2  | 100%  | 100%  | 100%  | 99.3% | Developmental and epileptic encephalopathy 32, 616366                                                                     |
| KCNC1  | 100%  | 100%  | 100%  | 98%   | Epilepsy, progressive myoclonic 7, 616187                                                                                 |
| KCNC3  | 99.6% | 97.6% | 98.9% | 87.9% | Spinocerebellar ataxia 13, 605259                                                                                         |
| KCND3  | 100%  | 100%  | 100%  | 97.8% | Spinocerebellar ataxia 19, 607346;Brugada syndrome 9, 616399                                                              |
| KCNJ10 | 100%  | 100%  | 100%  | 99.3% | Enlarged vestibular aqueduct, digenic, 600791;SESAME syndrome, 612780                                                     |

|           |      |      |      |       |                                                                                                                                                                                                                                                   |
|-----------|------|------|------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| KCNJ6     | 100% | 100% | 100% | 99.6% | Keppen-Lubinsky syndrome, 614098                                                                                                                                                                                                                  |
| KCNMA1    | 100% | 100% | 100% | 99%   | {Epilepsy, idiopathic generalized, susceptibility to, 16}, 618596;Paroxysmal nonkinesigenic dyskinesia, 3, with or without generalized epilepsy, 609446;Cerebellar atrophy, developmental delay, and seizures, 617643;Liang-Wang syndrome, 618729 |
| KCTD17    | 100% | 100% | 100% | 97.2% | Dystonia 26, myoclonic, 616398                                                                                                                                                                                                                    |
| KCTD7     | 100% | 100% | 100% | 99.1% | Epilepsy, progressive myoclonic 3, with or without intracellular inclusions, 611726                                                                                                                                                               |
| KIDINS220 | 100% | 100% | 100% | 99.6% | Spastic paraplegia, intellectual disability, nystagmus, and obesity, 617296;Ventriculomegaly and arthrogryposis, 619501                                                                                                                           |
| KIF1A     | 100% | 100% | 100% | 98.9% | NESCAV syndrome, 614255;Neuropathy, hereditary sensory, type IIC, 614213;Spastic paraplegia 30, autosomal dominant, 610357;Spastic paraplegia 30, autosomal recessive, 620607                                                                     |
| KIF1C     | 100% | 100% | 100% | 98.8% | Spastic ataxia 2, autosomal recessive, 611302                                                                                                                                                                                                     |

|       |       |       |       |       |                                                                                                                                                           |
|-------|-------|-------|-------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| KIF5A | 100%  | 100%  | 100%  | 99.1% | Myoclonus, intractable, neonatal, 617235;{Amyotrophic lateral sclerosis, susceptibility to, 25}, 617921;Spastic paraplegia 10, autosomal dominant, 604187 |
| KMT2B | 99.8% | 99.4% | 100%  | 98.2% | Intellectual developmental disorder, autosomal dominant 68, 619934;Dystonia 28, childhood-onset, 617284                                                   |
| KPNA3 | 100%  | 100%  | 100%  | 99.6% | Spastic paraplegia 88, autosomal dominant, 620106                                                                                                         |
| L1CAM | 100%  | 99.7% | 98.4% | 68.6% | MASA syndrome, 303350;Hydrocephalus, congenital, X-linked, 307000;?Corpus callosum, partial agenesis of, 304100                                           |
| LAMA1 | 100%  | 100%  | 100%  | 99.5% | Poretti-Boltshauser syndrome, 615960                                                                                                                      |
| LAMB1 | 100%  | 99.8% | 100%  | 99.7% | Lissencephaly 5, 615191                                                                                                                                   |
| LMNB1 | 100%  | 100%  | 100%  | 99.4% | Leukodystrophy, adult-onset, autosomal dominant, 169500;Microcephaly 26, primary, autosomal dominant, 619179                                              |
| MAG   | 100%  | 100%  | 100%  | 99%   | Spastic paraplegia 75, autosomal recessive, 616680                                                                                                        |

|          |      |       |       |       |                                                                                                                                                                                                                                                                                                                                         |
|----------|------|-------|-------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| MAPK8IP3 | 100% | 100%  | 100%  | 99.2% | Neurodevelopmental disorder with or without variable brain abnormalities, 618443                                                                                                                                                                                                                                                        |
| MARS2    | 100% | 100%  | 100%  | 99.8% | ?Combined oxidative phosphorylation deficiency 25, 616430;Spastic ataxia 3, autosomal recessive, 611390                                                                                                                                                                                                                                 |
| MBIP     | 100% | 100%  | 100%  | 99.7% |                                                                                                                                                                                                                                                                                                                                         |
| MECP2    | 100% | 99.3% | 97.1% | 64.9% | Rett syndrome, atypical, 312750;Encephalopathy, neonatal severe, 300673;Intellectual developmental disorder, X-linked syndromic, Lubs type, 300260;{Autism susceptibility, X-linked 3}, 300496;Intellectual developmental disorder, X-linked syndromic 13, 300055;Rett syndrome, 312750;Rett syndrome, preserved speech variant, 312750 |
| MECR     | 100% | 100%  | 100%  | 99.2% | Dystonia, childhood-onset, with optic atrophy and basal ganglia abnormalities, 617282;Optic atrophy 16, 620629                                                                                                                                                                                                                          |

|        |       |       |      |       |                                                                                                                                                               |
|--------|-------|-------|------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|
| MFF    | 95.9% | 95.9% | 100% | 99.7% | Encephalopathy due to defective mitochondrial and peroxisomal fission 2, 617086                                                                               |
| MFSD8  | 100%  | 100%  | 100% | 99.9% | Macular dystrophy with central cone involvement, 616170;Ceroid lipofuscinosis, neuronal, 7, 610951                                                            |
| MICU1  | 100%  | 100%  | 100% | 99.8% | Myopathy with extrapyramidal signs, 615673                                                                                                                    |
| MLC1   | 100%  | 100%  | 100% | 98.7% | Megalencephalic leukoencephalopathy with subcortical cysts 1, 604004                                                                                          |
| MMADHC | 89.3% | 89.3% | 100% | 99.3% | Methylmalonic aciduria and homocystinuria, cbID type, 277410;Methylmalonic aciduria, cbID type, 620953;Homocystinuria-megaloblastic anemia, cbID type, 620952 |
| MRE11  | 100%  | 100%  | 100% | 99.6% | Ataxia-telangiectasia-like disorder 1, 604391                                                                                                                 |



|        |      |       |      |       |                                                                                                                                                                                                                               |
|--------|------|-------|------|-------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| MTHFR  | 100% | 100%  | 100% | 99.1% | {Vascular disease, susceptibility to};Homocystinuria due to MTHFR deficiency, 236250;{Thromboembolism, susceptibility to}, 188050;{Schizophrenia, susceptibility to}, 181500;{Neural tube defects, susceptibility to}, 601634 |
| MTHFS  | 100% | 100%  | 100% | 98%   | Neurodevelopmental disorder with microcephaly, epilepsy, and hypomyelination, 618367                                                                                                                                          |
| MTPAP  | 100% | 100%  | 100% | 99.4% | ?Spastic ataxia 4, autosomal recessive, 613672                                                                                                                                                                                |
| MTRFR  | 100% | 99.8% | 100% | 98.8% | Spastic paraplegia 55, autosomal recessive, 615035;Combined oxidative phosphorylation deficiency 7, 613559                                                                                                                    |
| MTTP   | 100% | 100%  | 100% | 99.9% | Abetalipoproteinemia, 200100                                                                                                                                                                                                  |
| MYBPC1 | 100% | 100%  | 100% | 99.7% | Congenital myopathy 16, 618524;Lethal congenital contracture syndrome 4, 614915;Arthrogryposis, distal, type 1B, 614335                                                                                                       |
| MYORG  | 100% | 100%  | 100% | 98.8% | Basal ganglia calcification, idiopathic, 7, autosomal recessive, 618317                                                                                                                                                       |

|        |       |       |      |       |                                                                                                                                                        |
|--------|-------|-------|------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| NANS   | 100%  | 100%  | 100% | 99.1% | Spondyloepimetaphyseal dysplasia, Genevieve type, 610442                                                                                               |
| NARS2  | 92.3% | 92.3% | 100% | 99.7% | Combined oxidative phosphorylation deficiency 24, 616239;?Deafness, autosomal recessive 94, 618434                                                     |
| NDUFS7 | 100%  | 100%  | 100% | 98.3% | Mitochondrial complex I deficiency, nuclear type 3, 618224                                                                                             |
| NEFL   | 100%  | 100%  | 100% | 98.1% | Charcot-Marie-Tooth disease, type 1F, 607734;Charcot-Marie-Tooth disease, dominant intermediate G, 617882;Charcot-Marie-Tooth disease, type 2E, 607684 |
| NEU1   | 100%  | 100%  | 100% | 99.3% | Sialidosis, type II, 256550;Sialidosis, type I, 256550                                                                                                 |
| NEXMIF | 100%  | 100%  | 99%  | 72.1% | Intellectual developmental disorder, X-linked 98, 300912                                                                                               |
| NF2    | 100%  | 100%  | 100% | 99.3% | Meningioma, NF2-related, somatic, 607174;Schwannomatosis, vestibular, 101000;Schwannomatosis, somatic, 101000                                          |

|        |      |      |      |       |                                                                                                                                                        |
|--------|------|------|------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| NFU1   | 100% | 100% | 100% | 99.8% | Spastic paraplegia 93, autosomal recessive, 620938;Multiple mitochondrial dysfunctions syndrome 1, 605711                                              |
| NGLY1  | 100% | 100% | 100% | 99.3% | Congenital disorder of deglycosylation 1, 615273                                                                                                       |
| NIPA1  | 100% | 100% | 100% | 96.7% | Spastic paraplegia 6, autosomal dominant, 600363                                                                                                       |
| NKX2-1 | 100% | 100% | 100% | 98.7% | Chorea, hereditary benign, 118700;{Thyroid cancer, nonmedullary, 1}, 188550;Choreoathetosis, hypothyroidism, and neonatal respiratory distress, 610978 |
| NKX6-2 | 100% | 100% | 100% | 96.4% | Spastic ataxia 8, autosomal recessive, with hypomyelinating leukodystrophy, 617560                                                                     |
| NOL3   | 100% | 100% | 100% | 99.6% | ?Myoclonus, familial, 1, 614937                                                                                                                        |
| NPC1   | 100% | 100% | 100% | 99.4% | Niemann-Pick disease, type C1, 257220;Niemann-Pick disease, type D, 257220                                                                             |
| NPC2   | 100% | 100% | 100% | 99.1% | Niemann-pick disease, type C2, 607625                                                                                                                  |
| NPTX1  | 100% | 100% | 100% | 96.9% | Spinocerebellar ataxia 50, 620158                                                                                                                      |

|       |       |       |       |       |                                                                                                                                                                                                                          |
|-------|-------|-------|-------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NR4A2 | 100%  | 100%  | 100%  | 99.1% | Intellectual developmental disorder with language impairment and early-onset DOPA-responsive dystonia-parkinsonism, 619911                                                                                               |
| NT5C2 | 100%  | 100%  | 100%  | 99.6% | Spastic paraplegia 45, autosomal recessive, 613162                                                                                                                                                                       |
| NUP62 | 100%  | 100%  | 99.9% | 98.7% | Striatonigral degeneration, infantile, 271930                                                                                                                                                                            |
| NUS1  | 100%  | 100%  | 100%  | 99.1% | Intellectual developmental disorder, autosomal dominant 55, with seizures, 617831;?Congenital disorder of glycosylation, type 1aa, 617082                                                                                |
| OCLN  | 94.5% | 94.5% | 100%  | 98.4% | Pseudo-TORCH syndrome 1, 251290                                                                                                                                                                                          |
| OGDHL | 100%  | 100%  | 100%  | 99.4% | Yoon-Bellen neurodevelopmental syndrome, 619701                                                                                                                                                                          |
| OPA1  | 100%  | 100%  | 100%  | 99.6% | Optic atrophy plus syndrome, 125250;{Glaucoma, normal tension, susceptibility to}, 606657;Optic atrophy 1, 165500;Behr syndrome, 210000;?Mitochondrial DNA depletion syndrome 14 (encephalocardiomyopathic type), 616896 |

|        |       |       |       |       |                                                                                                                                                                                                                                                                                                                   |
|--------|-------|-------|-------|-------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| OPHN1  | 94%   | 93.8% | 99.3% | 72.4% | Intellectual developmental disorder, X-linked syndromic, Billuart type, 300486                                                                                                                                                                                                                                    |
| PACS2  | 100%  | 99.9% | 100%  | 98.4% | Developmental and epileptic encephalopathy 66, 618067                                                                                                                                                                                                                                                             |
| PANK2  | 100%  | 100%  | 100%  | 99.8% | Neurodegeneration with brain iron accumulation 1, 234200                                                                                                                                                                                                                                                          |
| PAX6   | 100%  | 100%  | 100%  | 99.1% | Optic nerve hypoplasia, 165550;Cataract with late-onset corneal dystrophy, 106210;Microphthalmia/coloboma 12, 120200;?Coloboma of optic nerve, 120430;Aniridia, 106210;Anterior segment dysgenesis 5, multiple subtypes, 604229;?Morning glory disc anomaly, 120430;Foveal hypoplasia 1, 136520;Keratitis, 148190 |
| PCYT2  | 100%  | 100%  | 100%  | 98.9% | Spastic paraplegia 82, autosomal recessive, 618770                                                                                                                                                                                                                                                                |
| PDE10A | 99.5% | 97.9% | 99.6% | 91.8% | Striatal degeneration, autosomal dominant, 616922;Dyskinesia, limb and orofacial, infantile-onset, 616921                                                                                                                                                                                                         |

|        |       |       |       |       |                                                                                                                                                                                                                                    |
|--------|-------|-------|-------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PDE2A  | 100%  | 100%  | 100%  | 99.2% | Intellectual developmental disorder with paroxysmal dyskinesia or seizures, 619150                                                                                                                                                 |
| PDE8B  | 100%  | 100%  | 100%  | 99.4% | Pigmented nodular adrenocortical disease, primary, 3, 614190;Striatal degeneration, autosomal dominant, 609161                                                                                                                     |
| PDGFB  | 100%  | 100%  | 100%  | 99.2% | Meningioma, SIS-related, 607174;Basal ganglia calcification, idiopathic, 5, 615483;Dermatofibrosarcoma protuberans, 607907                                                                                                         |
| PDGFRB | 100%  | 100%  | 100%  | 99%   | Premature aging syndrome, Penttinen type, 601812;Kosaki overgrowth syndrome, 616592;Myofibromatosis, infantile, 1, 228550;Basal ganglia calcification, idiopathic, 4, 615007;Myeloproliferative disorder with eosinophilia, 131440 |
| PDHA1  | 99.1% | 93.7% | 98.6% | 71.4% | Pyruvate dehydrogenase E1-alpha deficiency, 312170                                                                                                                                                                                 |
| PDHX   | 100%  | 100%  | 100%  | 99.3% | Lacticacidemia due to PDX1 deficiency, 245349                                                                                                                                                                                      |
| PDSS1  | 100%  | 100%  | 100%  | 99.5% | Coenzyme Q10 deficiency, primary, 2, 614651                                                                                                                                                                                        |

|        |       |       |      |       |                                                                                                                                   |
|--------|-------|-------|------|-------|-----------------------------------------------------------------------------------------------------------------------------------|
| PDSS2  | 100%  | 100%  | 100% | 99.3% | Coenzyme Q10 deficiency, primary, 3, 614652                                                                                       |
| PDYN   | 100%  | 100%  | 100% | 99.6% | Spinocerebellar ataxia 23, 610245                                                                                                 |
| PEX10  | 100%  | 100%  | 100% | 99.6% | Peroxisome biogenesis disorder 6A (Zellweger), 614870;Peroxisome biogenesis disorder 6B, 614871                                   |
| PEX2   | 100%  | 100%  | 100% | 99.7% | Peroxisome biogenesis disorder 5A (Zellweger), 614866;Peroxisome biogenesis disorder 5B, 614867                                   |
| PEX7   | 97.9% | 97.9% | 100% | 99.6% | Rhizomelic chondrodysplasia punctata, type 1, 215100;Peroxisome biogenesis disorder 9B, 614879                                    |
| PHYH   | 100%  | 100%  | 100% | 99.2% | Refsum disease, 266500                                                                                                            |
| PIGG   | 100%  | 100%  | 100% | 99.4% | [Blood group, EMM system], 619812;Neurodevelopmental disorder with or without hypotonia, seizures, and cerebellar atrophy, 616917 |
| PIK3R5 | 100%  | 100%  | 100% | 98.6% | Ataxia-oculomotor apraxia 3, 615217                                                                                               |

|        |      |       |       |       |                                                                                                                                                                                                                                                             |
|--------|------|-------|-------|-------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PLA2G6 | 100% | 99.7% | 100%  | 99.1% | Parkinson disease 14, autosomal recessive, 612953;Neurodegeneration with brain iron accumulation 2B, 610217;Infantile neuroaxonal dystrophy 1, 256600                                                                                                       |
| PLP1   | 100% | 99.6% | 99.1% | 71.8% | Pelizaeus-Merzbacher disease, 312080;Spastic paraplegia 2, X-linked, 312920                                                                                                                                                                                 |
| PMM2   | 100% | 100%  | 100%  | 99.7% | Congenital disorder of glycosylation, type Ia, 212065                                                                                                                                                                                                       |
| PMP22  | 100% | 100%  | 100%  | 99.5% | Charcot-Marie-Tooth disease, type 1A, 118220;Roussy-Levy syndrome, 180800;Charcot-Marie-Tooth disease, type 1E, 118300;?Neuropathy, inflammatory demyelinating, 139393;Neuropathy, recurrent, with pressure palsies, 162500;Dejerine-Sottas disease, 145900 |
| PMPCA  | 96%  | 96%   | 100%  | 99.1% | Spinocerebellar ataxia, autosomal recessive 2, 213200                                                                                                                                                                                                       |
| PNKD   | 100% | 100%  | 100%  | 98.7% | Paroxysmal nonkinesigenic dyskinesia 1, 118800                                                                                                                                                                                                              |



|        |      |      |      |       |                                                                                                                                                                                    |
|--------|------|------|------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PNKP   | 100% | 100% | 100% | 97.9% | ?Charcot-Marie-Tooth disease, type 2B2, 605589;Ataxia-oculomotor apraxia 4, 616267;Microcephaly, seizures, and developmental delay, 613402                                         |
| PNPLA6 | 100% | 100% | 100% | 98.8% | Spastic paraplegia 39, autosomal recessive, 612020;Oliver-McFarlane syndrome, 275400;?Laurence-Moon syndrome, 245800;Boucher-Neuhauser syndrome, 215470                            |
| PNPT1  | 100% | 100% | 100% | 99.8% | Spinocerebellar ataxia 25, 608703;Deafness, autosomal recessive 70, with or without adult-onset neurodegeneration, 614934;Combined oxidative phosphorylation deficiency 13, 614932 |

|        |       |       |      |       |                                                                                                                                                                                                                                                                                                                                             |
|--------|-------|-------|------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| POLG   | 100%  | 100%  | 100% | 99.5% | Mitochondrial recessive ataxia syndrome (includes SANDO and SCAE), 607459;Mitochondrial DNA depletion syndrome 4B (MNGIE type), 613662;Mitochondrial DNA depletion syndrome 4A (Alpers type), 203700;Progressive external ophthalmoplegia, autosomal dominant 1, 157640;Progressive external ophthalmoplegia, autosomal recessive 1, 258450 |
| POLR1C | 83.5% | 83.2% | 100% | 98.8% | Leukodystrophy, hypomyelinating, 11, 616494;Treacher Collins syndrome 3, 248390                                                                                                                                                                                                                                                             |
| POLR3A | 100%  | 100%  | 100% | 99.3% | Wiedemann-Rautenstrauch syndrome, 264090;Leukodystrophy, hypomyelinating, 7, with or without oligodontia and/or hypogonadotropic hypogonadism, 607694                                                                                                                                                                                       |

|          |       |       |       |       |                                                                                                                                                                          |
|----------|-------|-------|-------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| POLR3B   | 100%  | 100%  | 100%  | 99.8% | Leukodystrophy, hypomyelinating, 8, with or without oligodontia and/or hypogonadotropic hypogonadism, 614381;Charcot-Marie-Tooth disease, demyelinating, type 1I, 619742 |
| POU4F1   | 93%   | 86.6% | 99.9% | 91.4% | Ataxia, intention tremor, and hypotonia syndrome, childhood-onset, 619352                                                                                                |
| PPP1R3F  | 100%  | 99.9% | 97.9% | 67.5% |                                                                                                                                                                          |
| PPT1     | 90.3% | 90.3% | 100%  | 99.7% | Ceroid lipofuscinosis, neuronal, 1, 256730                                                                                                                               |
| PRDX3    | 100%  | 100%  | 100%  | 99.8% | Spinocerebellar ataxia, autosomal recessive 32, 619862;Corneal dystrophy, punctiform and polychromatic pre-Descemet, 619871                                              |
| PRF1     | 100%  | 100%  | 100%  | 99%   | Hemophagocytic lymphohistiocytosis, familial, 2, 603553;Aplastic anemia, 609135;Lymphoma, non-Hodgkin, 605027                                                            |
| PRICKLE1 | 100%  | 100%  | 100%  | 99.6% | Epilepsy, progressive myoclonic 1B, 612437                                                                                                                               |
| PRKCG    | 100%  | 100%  | 100%  | 97.5% | Spinocerebellar ataxia 14, 605361                                                                                                                                        |

|       |      |      |      |       |                                                                                                                                                                                                                                       |
|-------|------|------|------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PRKN  | 100% | 100% | 100% | 99.2% | Adenocarcinoma of lung, somatic, 211980;Parkinson disease, juvenile, type 2, 600116;Ovarian cancer, somatic, 167000                                                                                                                   |
| PRKRA | 100% | 100% | 100% | 99.8% | Dystonia 16, 612067                                                                                                                                                                                                                   |
| PRRT2 | 100% | 100% | 100% | 98.1% | Convulsions, familial infantile, with paroxysmal choreoathetosis, 602066;Seizures, benign familial infantile, 2, 605751;Episodic kinesigenic dyskinesia 1, 128200                                                                     |
| PSAP  | 100% | 100% | 100% | 99.3% | Combined SAP deficiency, 611721;Krabbe disease, atypical, 611722;Metachromatic leukodystrophy due to SAP-b deficiency, 249900;Gaucher disease, atypical, 610539;{Parkinson disease 24, autosomal dominant, susceptibility to}, 619491 |
| PTRH2 | 100% | 100% | 100% | 99.2% | Infantile-onset multisystem neurologic, endocrine, and pancreatic disease, 616263                                                                                                                                                     |
| PTS   | 100% | 100% | 100% | 99.2% | Hyperphenylalaninemia, BH4-deficient, A, 261640                                                                                                                                                                                       |

|          |       |       |       |       |                                                                                                                                  |
|----------|-------|-------|-------|-------|----------------------------------------------------------------------------------------------------------------------------------|
| PUM1     | 100%  | 100%  | 99.9% | 98.1% | Spinocerebellar ataxia 47, 617931;Neurodevelopmental disorder with motor abnormalities, seizures, and facial dysmorphism, 620719 |
| PYCR2    | 100%  | 100%  | 100%  | 99.4% | Leukodystrophy, hypomyelinating, 10, 616420                                                                                      |
| QDPR     | 100%  | 100%  | 100%  | 99.6% | Hyperphenylalaninemia, BH4-deficient, C, 261630                                                                                  |
| RAB18    | 100%  | 100%  | 100%  | 99.8% | Warburg micro syndrome 3, 614222                                                                                                 |
| RAB3GAP1 | 100%  | 100%  | 100%  | 99.7% | Martsolf syndrome 2, 619420;Warburg micro syndrome 1, 600118                                                                     |
| RAB3GAP2 | 94.4% | 94.4% | 100%  | 99.5% | Martsolf syndrome 1, 212720;Warburg micro syndrome 2, 614225                                                                     |
| RAD51    | 89.3% | 89.3% | 100%  | 99.4% | Mirror movements 2, 614508;{Breast cancer, susceptibility to}, 114480;Fanconi anemia, complementation group R, 617244            |
| RARS1    | 94.5% | 94.3% | 100%  | 99.3% | Leukodystrophy, hypomyelinating, 9, 616140                                                                                       |
| RARS2    | 94%   | 92.7% | 100%  | 99.7% | Pontocerebellar hypoplasia, type 6, 611523                                                                                       |

|          |       |       |      |       |                                                                                                                                                                                              |
|----------|-------|-------|------|-------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| REEP1    | 85.8% | 85.8% | 100% | 99.2% | Neuronopathy, distal hereditary motor, autosomal recessive 6, 620011;Spastic paraplegia 31, autosomal dominant, 610250;?Neuronopathy, distal hereditary motor, autosomal dominant 12, 614751 |
| REEP2    | 100%  | 100%  | 100% | 98.2% | Spastic paraplegia 72A, autosomal dominant, 615625;?Spastic paraplegia 72B, autosomal recessive, 620606                                                                                      |
| RFC1     | 100%  | 100%  | 100% | 99.6% | Cerebellar ataxia, neuropathy, and vestibular areflexia syndrome, 614575                                                                                                                     |
| RHOBTB2  | 98.7% | 98.7% | 100% | 99.2% | Developmental and epileptic encephalopathy 64, 618004                                                                                                                                        |
| RNASEH2A | 100%  | 100%  | 100% | 99.2% | Aicardi-Goutieres syndrome 4, 610333                                                                                                                                                         |
| RNASEH2B | 91.4% | 91.4% | 100% | 99.2% | Aicardi-Goutieres syndrome 2, 610181                                                                                                                                                         |
| RNASEH2C | 100%  | 100%  | 100% | 98.1% | Aicardi-Goutieres syndrome 3, 610329                                                                                                                                                         |
| RNF170   | 100%  | 100%  | 100% | 99.7% | Ataxia, sensory, 1, autosomal dominant, 608984;Spastic paraplegia 85, autosomal recessive, 619686                                                                                            |

|        |      |      |      |       |                                                                                                                                          |
|--------|------|------|------|-------|------------------------------------------------------------------------------------------------------------------------------------------|
| RNF216 | 100% | 100% | 100% | 99.8% | Cerebellar ataxia and hypogonadotropic hypogonadism, 212840                                                                              |
| RORA   | 100% | 100% | 100% | 98.4% | Intellectual developmental disorder with or without epilepsy or cerebellar ataxia, 618060                                                |
| RTN2   | 100% | 100% | 100% | 98.9% | Neuronopathy, distal hereditary motor, autosomal recessive 11, with spasticity, 620854;Spastic paraplegia 12, autosomal dominant, 604805 |
| RUBCN  | 100% | 100% | 100% | 99.1% | Spinocerebellar ataxia, autosomal recessive 15, 615705                                                                                   |
| SACS   | 99%  | 99%  | 100% | 99.8% | Spastic ataxia, Charlevoix-Saguenay type, 270550                                                                                         |
| SAMD9L | 100% | 100% | 100% | 99.8% | Ataxia-pancytopenia syndrome, 159550;?Spinocerebellar ataxia 49, 619806;Monosomy 7 myelodysplasia and leukemia syndrome 1, 252270        |
| SAMHD1 | 100% | 100% | 100% | 99.8% | ?Chilblain lupus 2, 614415;Aicardi-Goutieres syndrome 5, 612952                                                                          |

|        |      |       |      |       |                                                                                                                                                                                                                                           |
|--------|------|-------|------|-------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| SCN11A | 100% | 99.9% | 100% | 99.3% | Episodic pain syndrome, familial, 3, 615552;Neuropathy, hereditary sensory and autonomic, type VII, 615548                                                                                                                                |
| SCN1A  | 100% | 100%  | 100% | 99.7% | Developmental and epileptic encephalopathy 6B, non-Dravet, 619317;Migraine, familial hemiplegic, 3, 609634;Dravet syndrome, 607208;Febrile seizures, familial, 3A, 604403;Generalized epilepsy with febrile seizures plus, type 2, 604403 |
| SCN2A  | 100% | 100%  | 100% | 99.8% | Seizures, benign familial infantile, 3, 607745;Developmental and epileptic encephalopathy 11, 613721;Episodic ataxia, type 9, 618924                                                                                                      |
| SCN8A  | 100% | 100%  | 100% | 99.2% | ?Myoclonus, familial, 2, 618364;Seizures, benign familial infantile, 5, 617080;Cognitive impairment with or without cerebellar ataxia, 614306;Developmental and epileptic encephalopathy 13, 614558                                       |
| SCYL1  | 100% | 100%  | 100% | 98.6% | Spinocerebellar ataxia, autosomal recessive 21, 616719                                                                                                                                                                                    |



|         |       |       |      |       |                                                                                                                                  |
|---------|-------|-------|------|-------|----------------------------------------------------------------------------------------------------------------------------------|
| SELENOI | 100%  | 100%  | 100% | 99.9% | Spastic paraplegia 81, autosomal recessive, 618768                                                                               |
| SEPSECS | 99.4% | 95.2% | 100% | 99.4% | Pontocerebellar hypoplasia type 2D, 613811                                                                                       |
| SERAC1  | 100%  | 100%  | 100% | 99.8% | 3-methylglutaconic aciduria with deafness, encephalopathy, and Leigh-like syndrome, 614739                                       |
| SETX    | 100%  | 100%  | 100% | 99.5% | Spinocerebellar ataxia, autosomal recessive, with axonal neuropathy 2, 606002; Amyotrophic lateral sclerosis 4, juvenile, 602433 |
| SGCE    | 90.8% | 90.8% | 100% | 99.6% | Dystonia-11, myoclonic, 159900                                                                                                   |
| SIL1    | 100%  | 100%  | 100% | 99.5% | Marinesco-Sjogren syndrome, 248800                                                                                               |
| SLC12A6 | 100%  | 99.9% | 100% | 99.7% | Agenesis of the corpus callosum with peripheral neuropathy, 218000; Charcot-Marie-Tooth disease, axonal, type 2II, 620068        |
| SLC16A2 | 100%  | 99.8% | 99%  | 70.5% | Allan-Herndon-Dudley syndrome, 300523                                                                                            |
| SLC18A2 | 100%  | 100%  | 100% | 99.5% | Parkinsonism-dystonia, infantile, 2, 618049                                                                                      |

|          |       |       |      |       |                                                                                                                                                                                                                                                                            |
|----------|-------|-------|------|-------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| SLC19A3  | 99.8% | 98.7% | 100% | 99.6% | Thiamine metabolism dysfunction syndrome 2 (biotin/thiamine-responsive basal ganglia disease type), 607483                                                                                                                                                                 |
| SLC1A3   | 97.1% | 94.1% | 100% | 99.8% | Episodic ataxia, type 6, 612656                                                                                                                                                                                                                                            |
| SLC20A2  | 100%  | 100%  | 100% | 99.3% | Basal ganglia calcification, idiopathic, 1, 213600                                                                                                                                                                                                                         |
| SLC25A15 | 100%  | 100%  | 100% | 99.5% | Hyperornithinemia-hyperammonemia-homocitrullinemia syndrome, 238970                                                                                                                                                                                                        |
| SLC25A46 | 100%  | 100%  | 100% | 99.6% | Neuropathy, hereditary motor and sensory, type VIB, 616505;Pontocerebellar hypoplasia, type 1E, 619303                                                                                                                                                                     |
| SLC2A1   | 100%  | 100%  | 100% | 99.2% | Dystonia 9, 601042;GLUT1 deficiency syndrome 1, infantile onset, severe, 606777;Stomatin-deficient cryohydrocytosis with neurologic defects, 608885;{Epilepsy, idiopathic generalized, susceptibility to, 12}, 614847;GLUT1 deficiency syndrome 2, childhood onset, 612126 |
| SLC30A10 | 100%  | 100%  | 100% | 98.8% | Hypermanganesemia with dystonia 1, 613280                                                                                                                                                                                                                                  |

|          |       |       |      |       |                                                                                               |
|----------|-------|-------|------|-------|-----------------------------------------------------------------------------------------------|
| SLC33A1  | 100%  | 100%  | 100% | 99.4% | Spastic paraplegia 42, autosomal dominant, 612539;Huppke-Brendel syndrome, 614482             |
| SLC39A14 | 93.6% | 93.6% | 100% | 99.4% | ?Hyperostosis cranialis interna, 144755;Hypermanganesemia with dystonia 2, 617013             |
| SLC52A2  | 100%  | 100%  | 100% | 99.6% | Brown-Vialetto-Van Laere syndrome 2, 614707                                                   |
| SLC52A3  | 100%  | 100%  | 100% | 99.1% | ?Fazio-Londe disease, 211500;Brown-Vialetto-Van Laere syndrome 1, 211530                      |
| SLC6A3   | 100%  | 100%  | 100% | 99.2% | Parkinsonism-dystonia, infantile, 1, 613135;{Nicotine dependence, protection against}, 188890 |
| SLC6A5   | 100%  | 100%  | 100% | 98.7% | Hyperekplexia 3, 614618                                                                       |
| SLC9A1   | 100%  | 100%  | 100% | 99.1% | Lichtenstein-Knorr syndrome, 616291                                                           |
| SMDT1    | 100%  | 100%  | 100% | 99.4% |                                                                                               |
| SMPD1    | 100%  | 100%  | 100% | 99.1% | Niemann-Pick disease, type B, 607616;Niemann-Pick disease, type A, 257200                     |
| SNORD118 |       |       |      |       | Leukoencephalopathy, brain calcifications, and cysts, 614561                                  |
| SNX14    | 95%   | 95%   | 100% | 99.9% | Spinocerebellar ataxia, autosomal recessive 20, 616354                                        |

|       |       |       |      |       |                                                                                                                                                           |
|-------|-------|-------|------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| SOX10 | 97.8% | 97.8% | 100% | 98.3% | Waardenburg syndrome, type 4C, 613266;PCWH syndrome, 609136;Waardenburg syndrome, type 2E, with or without neurologic involvement, 611584                 |
| SPART | 100%  | 100%  | 100% | 99.7% | Troyer syndrome, 275900                                                                                                                                   |
| SPAST | 100%  | 99.9% | 100% | 98.5% | Spastic paraplegia 4, autosomal dominant, 182601                                                                                                          |
| SPG11 | 99.6% | 99.6% | 100% | 99.5% | Amyotrophic lateral sclerosis 5, juvenile, 602099;Charcot-Marie-Tooth disease, axonal, type 2X, 616668;Spastic paraplegia 11, autosomal recessive, 604360 |
| SPG21 | 100%  | 100%  | 100% | 99.7% | Mast syndrome, 248900                                                                                                                                     |
| SPG7  | 100%  | 100%  | 100% | 98.7% | Spastic paraplegia 7, autosomal recessive, 607259                                                                                                         |
| SPR   | 100%  | 100%  | 100% | 98.9% | Dystonia, dopa-responsive, due to sepiapterin reductase deficiency, 612716                                                                                |

|        |       |       |      |       |                                                                                                                                                                                                                                                                     |
|--------|-------|-------|------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| SPTAN1 | 99.3% | 98.9% | 100% | 99.4% | Developmental delay with or without epilepsy, 620540;Developmental and epileptic encephalopathy 5, 613477;Spastic paraplegia 91, autosomal dominant, with or without cerebellar ataxia, 620538;Neuronopathy, distal hereditary motor, autosomal dominant 11, 620528 |
| SPTBN2 | 100%  | 100%  | 100% | 99%   | Spinocerebellar ataxia 5, 600224;Spinocerebellar ataxia, autosomal recessive 14, 615386                                                                                                                                                                             |
| SQSTM1 | 100%  | 100%  | 100% | 98.9% | Neurodegeneration with ataxia, dystonia, and gaze palsy, childhood-onset, 617145;Frontotemporal dementia and/or amyotrophic lateral sclerosis 3, 616437;Myopathy, distal, with rimmed vacuoles, 617158;Paget disease of bone 3, 167250                              |
| STUB1  | 100%  | 100%  | 100% | 97.8% | Spinocerebellar ataxia 48, 618093;Spinocerebellar ataxia, autosomal recessive 16, 615768                                                                                                                                                                            |
| SUMF1  | 100%  | 100%  | 100% | 99.5% | Multiple sulfatase deficiency, 272200                                                                                                                                                                                                                               |

|         |       |       |      |       |                                                                                                                                                                                   |
|---------|-------|-------|------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| SUOX    | 100%  | 100%  | 100% | 98.7% | Sulfite oxidase deficiency, 272300                                                                                                                                                |
| SYNE1   | 100%  | 100%  | 100% | 99.6% | Arthrogryposis multiplex congenita 3, myogenic type, 618484;Emery-Dreifuss muscular dystrophy 4, autosomal dominant, 612998;Spinocerebellar ataxia, autosomal recessive 8, 610743 |
| TANGO2  | 100%  | 100%  | 100% | 99.5% | Metabolic encephalomyopathic crises, recurrent, with rhabdomyolysis, cardiac arrhythmias, and neurodegeneration, 616878                                                           |
| TBC1D20 | 92.2% | 92.2% | 100% | 99.5% | Warburg micro syndrome 4, 615663                                                                                                                                                  |
| TBC1D23 | 100%  | 100%  | 100% | 99.8% | Pontocerebellar hypoplasia, type 11, 617695                                                                                                                                       |
| TBCD    | 91.2% | 90.5% | 100% | 99.2% | Encephalopathy, progressive, early-onset, with brain atrophy and thin corpus callosum, 617193                                                                                     |
| TDP1    | 100%  | 100%  | 100% | 99.8% | ?Spinocerebellar ataxia, autosomal recessive, with axonal neuropathy 1, 607250                                                                                                    |
| TDP2    | 100%  | 100%  | 100% | 99.7% | Spinocerebellar ataxia, autosomal recessive 23, 616949                                                                                                                            |

|          |       |       |       |       |                                                                                                                                                                                   |
|----------|-------|-------|-------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| TECPR2   | 100%  | 100%  | 100%  | 99.4% | Neuropathy, hereditary sensory and autonomic, type IX, with developmental delay, 615031                                                                                           |
| TENM4    | 100%  | 100%  | 100%  | 99.3% | Essential tremor, hereditary, 5, 616736                                                                                                                                           |
| TGM6     | 100%  | 100%  | 100%  | 99.4% | Spinocerebellar ataxia 35, 613908                                                                                                                                                 |
| TH       | 100%  | 100%  | 100%  | 99.1% | Segawa syndrome, recessive, 605407                                                                                                                                                |
| THAP1    | 100%  | 100%  | 100%  | 99.6% | Dystonia 6, torsion, 602629                                                                                                                                                       |
| TIMM8A   | 100%  | 99.5% | 99.2% | 71%   | Mohr-Tranebjaerg syndrome, 304700                                                                                                                                                 |
| TMEM106B | 100%  | 100%  | 100%  | 99.8% | Leukodystrophy, hypomyelinating, 16, 617964                                                                                                                                       |
| TMEM240  | 100%  | 100%  | 100%  | 95.1% | Spinocerebellar ataxia 21, 607454                                                                                                                                                 |
| TMEM67   | 96.1% | 96.1% | 100%  | 99.8% | Nephronophthisis 11, 613550;{Bardet-Biedl syndrome 14, modifier of}, 615991;Joubert syndrome 6, 610688;Meckel syndrome 3, 607361;?RHYNS syndrome, 602152;COACH syndrome 1, 216360 |
| TOE1     | 100%  | 100%  | 100%  | 99.6% | Pontocerebellar hypoplasia, type 7, 614969                                                                                                                                        |

|       |       |       |      |       |                                                                                                                                                                                                                                             |
|-------|-------|-------|------|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| TOR1A | 93.4% | 90.8% | 100% | 98.8% | {Dystonia-1, modifier of};Arthrogryposis multiplex congenita 5, 618947;Dystonia-1, torsion, 128100                                                                                                                                          |
| TPP1  | 100%  | 100%  | 100% | 99.2% | Ceroid lipofuscinosis, neuronal, 2, 204500;Spinocerebellar ataxia, autosomal recessive 7, 609270                                                                                                                                            |
| TREM2 | 100%  | 100%  | 100% | 99.5% | {Alzheimer disease 17, susceptibility to}, 615080;Polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy 2, 618193                                                                                                    |
| TREX1 | 100%  | 100%  | 100% | 99.7% | Vasculopathy, retinal, with cerebral leukoencephalopathy and systemic manifestations, 192315;Aicardi-Goutieres syndrome 1, dominant and recessive, 225750;{Systemic lupus erythematosus, susceptibility to}, 152700;Chilblain lupus, 610448 |
| TRIT1 | 100%  | 100%  | 100% | 99.5% | Combined oxidative phosphorylation deficiency 35, 617873                                                                                                                                                                                    |



|         |       |       |      |       |                                                                                                                                                                       |
|---------|-------|-------|------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| TRPM3   | 97.8% | 97.8% | 100% | 99.6% | ?Cataract 50 with or without glaucoma, 620253;Neurodevelopmental disorder with hypotonia, dysmorphic facies, and skeletal anomalies, with or without seizures, 620224 |
| TSEN15  | 100%  | 100%  | 100% | 99.5% | Pontocerebellar hypoplasia, type 2F, 617026                                                                                                                           |
| TSEN2   | 88.4% | 88.4% | 100% | 99.1% | Pontocerebellar hypoplasia type 2B, 612389                                                                                                                            |
| TSEN54  | 100%  | 100%  | 100% | 99%   | Pontocerebellar hypoplasia type 2A, 277470;Pontocerebellar hypoplasia type 4, 225753;?Pontocerebellar hypoplasia type 5, 610204                                       |
| TSPOAP1 | 100%  | 100%  | 100% | 98.7% | Dystonia 22, juvenile-onset, 620453;?Dystonia 22, adult-onset, 620456                                                                                                 |
| TTBK2   | 100%  | 100%  | 100% | 99.6% | Spinocerebellar ataxia 11, 604432                                                                                                                                     |
| TTC19   | 100%  | 100%  | 100% | 99.6% | Mitochondrial complex III deficiency, nuclear type 2, 615157                                                                                                          |
| TTPA    | 100%  | 100%  | 100% | 99%   | Ataxia with isolated vitamin E deficiency, 277460                                                                                                                     |
| TUBA1A  | 100%  | 100%  | 100% | 99.1% | Lissencephaly 3, 611603                                                                                                                                               |

|        |       |       |      |       |                                                                                                                                                                                                      |
|--------|-------|-------|------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| TUBB   | 99.8% | 99%   | 100% | 99.2% | Symmetric circumferential skin creases, congenital, 1, 156610;Cortical dysplasia, complex, with other brain malformations 6, 615771                                                                  |
| TUBB4A | 99.4% | 96.9% | 100% | 98.7% | Dystonia 4, torsion, autosomal dominant, 128101;Leukodystrophy, hypomyelinating, 6, 612438                                                                                                           |
| TUBG1  | 100%  | 100%  | 100% | 99%   | Cortical dysplasia, complex, with other brain malformations 4, 615412                                                                                                                                |
| TWINK  | 100%  | 100%  | 100% | 99.5% | Mitochondrial DNA depletion syndrome 7 (hepatocerebral type), 271245;Progressive external ophthalmoplegia with mitochondrial DNA deletions, autosomal dominant 3, 609286;Perrault syndrome 5, 616138 |
| TYROBP | 100%  | 100%  | 100% | 98.9% | Polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy 1, 221770                                                                                                               |
| UBA5   | 99.6% | 97.2% | 100% | 99.9% | ?Spinocerebellar ataxia, autosomal recessive 24, 617133;Developmental and epileptic encephalopathy 44, 617132                                                                                        |

|       |      |      |      |       |                                                                                                                                                                                                                  |
|-------|------|------|------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UBAP1 | 100% | 100% | 100% | 98.7% | Spastic paraplegia 80, autosomal dominant, 618418                                                                                                                                                                |
| UBTF  | 100% | 100% | 100% | 98.8% | Neurodegeneration, childhood-onset, with brain atrophy, 617672                                                                                                                                                   |
| UCHL1 | 100% | 100% | 100% | 99.7% | {?Parkinson disease 5, susceptibility to}, 613643;Spastic paraplegia 79A, autosomal dominant, 620221;Spastic paraplegia 79B, autosomal recessive, 615491                                                         |
| VAC14 | 100% | 100% | 100% | 99.4% | Striatonigral degeneration, childhood-onset, 617054                                                                                                                                                              |
| VAMP1 | 100% | 100% | 100% | 99.8% | Myasthenic syndrome, congenital, 25, 618323;Spastic ataxia 1, autosomal dominant, 108600                                                                                                                         |
| VAR2  | 100% | 100% | 100% | 98.6% | Combined oxidative phosphorylation deficiency 20, 615917                                                                                                                                                         |
| VCP   | 100% | 100% | 100% | 99.6% | Frontotemporal dementia and/or amyotrophic lateral sclerosis 6, 613954;Charcot-Marie-Tooth disease, type 2Y, 616687;Inclusion body myopathy with early-onset Paget disease and frontotemporal dementia 1, 167320 |

|        |       |       |      |       |                                                                                                                  |
|--------|-------|-------|------|-------|------------------------------------------------------------------------------------------------------------------|
| VLDLR  | 100%  | 100%  | 100% | 99.4% | Cerebellar hypoplasia, impaired intellectual development, and dysequilibrium syndrome 1, 224050                  |
| VPS11  | 100%  | 100%  | 100% | 99.2% | ?Dystonia 32, 619637;Leukodystrophy, hypomyelinating, 12, 616683                                                 |
| VPS13A | 100%  | 100%  | 100% | 99.8% | Choreoacanthocytosis, 200150                                                                                     |
| VPS13D | 100%  | 100%  | 100% | 99.7% | Spinocerebellar ataxia, autosomal recessive 4, 607317                                                            |
| VPS16  | 100%  | 100%  | 100% | 99.3% | Dystonia 30, 619291                                                                                              |
| VPS37A | 100%  | 100%  | 100% | 99.8% | Spastic paraplegia 53, autosomal recessive, 614898                                                               |
| VPS41  | 100%  | 100%  | 100% | 99.7% | Spinocerebellar ataxia, autosomal recessive 29, 619389                                                           |
| VPS53  | 83%   | 81.7% | 100% | 99.4% | Pontocerebellar hypoplasia, type 2E, 615851                                                                      |
| VRK1   | 99.3% | 97.7% | 100% | 100%  | Pontocerebellar hypoplasia type 1A, 607596;Neuronopathy, distal hereditary motor, autosomal recessive 10, 620542 |

|        |       |       |       |       |                                                                                                                                                                                                                    |
|--------|-------|-------|-------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| WASHC5 | 100%  | 100%  | 100%  | 99.7% | Ritscher-Schinzel syndrome 1, 220210;Spastic paraplegia 8, autosomal dominant, 603563                                                                                                                              |
| WDR26  | 93.8% | 93.8% | 100%  | 99.2% | Skraban-Deardorff syndrome, 617616                                                                                                                                                                                 |
| WDR45  | 100%  | 99.6% | 98.1% | 67.7% | Neurodegeneration with brain iron accumulation 5, 300894                                                                                                                                                           |
| WDR73  | 100%  | 100%  | 100%  | 99.2% | Galloway-Mowat syndrome 1, 251300                                                                                                                                                                                  |
| WDR81  | 100%  | 100%  | 100%  | 99.2% | Cerebellar ataxia, impaired intellectual development, and dysquilibrium syndrome 2, 610185;Hydrocephalus, congenital, 3, with brain anomalies, 617967                                                              |
| WFS1   | 91.2% | 91.2% | 100%  | 99.1% | Deafness, autosomal dominant 6/14/38, 600965;?Cataract 41, 116400;Wolfram-like syndrome, autosomal dominant, 614296;{Diabetes mellitus, noninsulin-dependent, association with}, 125853;Wolfram syndrome 1, 222300 |

|         |      |       |       |       |                                                                                                                                                                  |
|---------|------|-------|-------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| WVOX    | 100% | 100%  | 100%  | 99.5% | Esophageal squamous cell carcinoma, somatic, 133239;Developmental and epileptic encephalopathy 28, 616211;Spinocerebellar ataxia, autosomal recessive 12, 614322 |
| XK      | 100% | 98.9% | 99.2% | 70.9% | McLeod syndrome, 300842                                                                                                                                          |
| XPA     | 100% | 100%  | 100%  | 99.9% | Xeroderma pigmentosum, group A, 278700                                                                                                                           |
| XPR1    | 100% | 100%  | 100%  | 99.7% | Basal ganglia calcification, idiopathic, 6, 616413                                                                                                               |
| XRCC1   | 100% | 100%  | 100%  | 98.7% | ?Spinocerebellar ataxia, autosomal recessive 26, 617633                                                                                                          |
| ZC4H2   | 100% | 99.8% | 99.1% | 71.6% | Wieacker-Wolff syndrome, 314580;Wieacker-Wolff syndrome, female-restricted, 301041                                                                               |
| ZFYVE26 | 100% | 100%  | 100%  | 99.5% | Spastic paraplegia 15, autosomal recessive, 270700                                                                                                               |
| ZFYVE27 | 100% | 100%  | 100%  | 99.1% |                                                                                                                                                                  |

Gene symbols used follow HGCN guidelines: Gray KA, Yates B, Seal RL, Wright MW, Bruford EA. Nucleic Acids Res. 2015 Jan 43(Database issue):D1079-85.

TWIST X2 Covered 10x describes the percentage of a gene's coding sequence that is covered at least 10x when analyzed by WES using TWIST X2 chemistry.

TWIST X2 Covered 20x describes the percentage of a gene's coding sequence that is covered at least 20x when analyzed by WES using TWIST X2 chemistry.

srWGS GRCh38 Covered 10x describes the percentage of a gene's coding sequence that is covered at least 10x when analyzed by WGS mapped against GRCh38.

srWGS GRCh38 Covered 20x describes the percentage of a gene's coding sequence that is covered at least 20x when analyzed by WGS mapped against GRCh38.

non-protein coding genes are covered, but as coverage statistics are based on protein coding regions, statistics could not be generated.

OMIM release used for OMIM disease identifiers and descriptions : March 17th, 2023.

This list is accurate for panel version DG 4.0.0

*Ad 1. "No OMIM phenotype" signifies a gene without a current OMIM association Ad 2. OMIM phenotype descriptions between {} signify risk factors*