

COMPLEMENT MEDIATED RENAL DISEASE PANEL DG-4.4.0

(25 GENES)

<i>Gene</i>	<i>Twist X2 covered 10x</i>	<i>Twist X2 covered 20x</i>	<i>srWGS covered 10x</i>	<i>srWGS covered 15x</i>	<i>srWGS covered 20x</i>	<i>Associated Phenotype description and OMIM disease ID</i>
ADAMTS13	100%	100%	100%	100%	99.1%	Thrombotic thrombocytopenic purpura, hereditary, 274150
C2	100%	100%	100%	100%	99.6%	C2 deficiency, 217000;{Macular degeneration, age-related, 14, reduced risk of}, 615489
C3	97.5%	97.5%	100%	100%	98.7%	C3 deficiency, 613779;{Hemolytic uremic syndrome, atypical, susceptibility to, 5}, 612925;{Macular degeneration, age-related, 9}, 611378
C5	100%	100%	100%	100%	99.8%	C5 deficiency, 609536;[Eculizumab, poor response to], 615749

CD46	100%	100%	100%	100%	99.9%	{Hemolytic uremic syndrome, atypical, susceptibility to, 2}, 612922
CD55	97.5%	93.5%	100%	100%	99.7%	[Blood group Cromer], 613793;Complement hyperactivation, angiopathic thrombosis, and protein-losing enteropathy, 226300
CD59	100%	100%	100%	100%	99.8%	Hemolytic anemia, CD59-mediated, with or without immune-mediated polyneuropathy, 612300
CFB	100%	100%	100%	100%	99.3%	Complement factor B deficiency, 615561;{Hemolytic uremic syndrome, atypical, susceptibility to, 4}, 612924;{Macular degeneration, age-related, 14, reduced risk of}, 615489

CFH	97.5%	97.5%	100%	100%	99.7%	{Macular degeneration, age-related, 4}, 610698;Basal laminar drusen, 126700;Complement factor H deficiency, 609814;{Hemolytic uremic syndrome, atypical, susceptibility to, 1}, 235400
CFHR1	95.2%	94.6%	96%	89.8%	78.1%	{Macular degeneration, age-related, reduced risk of}, 603075;{Hemolytic uremic syndrome, atypical, susceptibility to}, 235400
CFHR2	76.4%	76.4%	100%	100%	99.7%	
CFHR3	94.7%	94.3%	97.3%	93%	83%	{Macular degeneration, age-related, reduced risk of}, 603075;{Hemolytic uremic syndrome, atypical, susceptibility to}, 235400
CFHR4	100%	100%	100%	99.7%	97.5%	
CFHR5	100%	100%	100%	100%	99.7%	Nephropathy due to CFHR5 deficiency, 614809

CFI	100%	100%	100%	100%	99.7%	{Hemolytic uremic syndrome, atypical, susceptibility to, 3}, 612923;{Macular degeneration, age-related, 13, susceptibility to}, 615439;Complement factor I deficiency, 610984
CFP	100%	100%	98.6%	90%	71.8%	Properdin deficiency, X-linked, 312060
DGKE	100%	100%	100%	100%	99.6%	{Hemolytic uremic syndrome, atypical, susceptibility to, 7}, 615008;Nephrotic syndrome, type 7, 615008
EXOSC3	100%	100%	100%	100%	99.3%	Pontocerebellar hypoplasia, type 1B, 614678
EXOSC5	100%	100%	100%	99.8%	98.8%	Cerebellar ataxia, brain abnormalities, and cardiac conduction defects, 619576
HSD11B2	100%	100%	100%	99.8%	98.4%	Apparent mineralocorticoid excess, 218030
IQGAP1	100%	100%	100%	99.8%	98.8%	
MMACHC	100%	100%	100%	100%	99.2%	Methylmalonic aciduria and homocystinuria, cblC type, 277400

PLG	100%	100%	100%	100%	99.7%	Dysplasminogenemia, 217090;Angioedema, hereditary, 4, 619360;Plasminogen deficiency, type I, 217090
THBD	100%	100%	100%	99.7%	97.9%	Thrombophilia 12 due to thrombomodulin defect, 614486;{Hemolytic uremic syndrome, atypical, susceptibility to, 6}, 612926
VTN	100%	100%	100%	100%	99.1%	

Gene symbols used follow HGNC guidelines: Gray KA, Yates B, Seal RL, Wright MW, Bruford EA. *Nucleic Acids Res.* 2015 Jan 43(Database issue):D1079-85.

TWIST X2 covered 10x describes the percentage of a gene's coding sequence that is covered at least 10x when analyzed by WES using TWIST X2 chemistry mapped against GRCh38.

TWIST X2 covered 20x describes the percentage of a gene's coding sequence that is covered at least 20x when analyzed by WES using TWIST X2 chemistry mapped against GRCh38.

srWGS covered 10x describes the percentage of a gene's coding sequence that is covered at least 10x when analyzed by WGS mapped against GRCh38.

srWGS covered 15x describes the percentage of a gene's coding sequence that is covered at least 15x when analyzed by WGS mapped against GRCh38.

srWGS covered 20x describes the percentage of a gene's coding sequence that is covered at least 20x when analyzed by WGS mapped against GRCh38.

non-protein coding genes are covered, but as coverage statistics are based on protein coding regions, statistics could not be generated.

OMIM release used for OMIM disease identifiers and descriptions : November 25th, 2024.

This list is accurate for panel version DG 4.3.0

Ad 1. Blank field signifies a gene without a current OMIM association Ad 2. OMIM phenotype descriptions between {} signify risk factors